Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis

Ronald P. Mensink



Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis

Ronald P. Mensink

Faculty of Health, Medicine and Life Sciences, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University, the Netherlands



WHO Library Cataloguing-in-Publication Data

Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis.

1. Fatty Acids. 2. Cholesterol, HDL. 3. Lipoproteins. 4. Dietary Fats. 5. Review. I. Mensink, Ronald P. II. World Health Organization.

ISBN 978 92 4 156534 9

(NLM classification: QU 90)

© World Health Organization 2016

All rights reserved. Publications of the World Health Organization are available on the WHO website (http://www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; email: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications – whether for sale or for non-commercial distribution – should be addressed to WHO Press through the WHO website (http://www.who.int/about/licensing/copyright_form/index. html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

 $The \, named \, authors \, alone \, are \, responsible \, for \, the \, views \, expressed \, in \, this \, publication.$

Design and layout: minimum graphics

Printed by the WHO Document Production Services, Geneva, Switzerland

Suggested citation: Mensink, RP. Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis. Geneva: World Health Organization; 2016.

Contents

Ac	know	ledgements	V
Αb	brevi	ations and acronyms	vi
1	Intro	oduction	1
	1.1	Background	1
	1.2	Objectives	1
2	Metl	hods	2
	2.1	Criteria for selecting studies to include in this review	2
		2.1.1 Study characteristics	2
		2.1.2 Outcomes	2
	2.2	Data collection and analysis	3
		2.2.1 Identification of studies	3
		2.2.2 Data extraction and management	3
		2.2.3 Assessment of risk of bias in included studies	3
		2.2.4 Analytical methods	5
3	Resi	ults	7
	3.1	Search results	7
	3.2	Included studies	7
	3.3	Effects of interventions	7
		3.3.1 Analysis of total SFA intake	7
		3.3.2 Subgroup and sensitivity analyses for total SFA	9
		3.3.3 Analysis of individual SFA	11
		3.3.4 Subgroup and sensitivity analyses for individual SFA	12
	3.4	Quality of the evidence	12
4	Stre	ngths and limitations of review	13
	4.1	Strengths	13
	4.2	Limitations	13
5	Con	clusion	14
6	Tabl	es	15
7	Figu	res	31
Ar	nex 1	. PICO questions	32
Ar	nex 2	. Priority outcomes	33
Ar	inex 3	. Search strategy	34
Ar	nex 4	. Residuals analysis	35

Annex 5.	Relationship between observed and predicted serum LDL cholesterol concentrations	36
Annex 6.	Results of <i>cis</i> -PUFA replacement	37
Annex 7.	Results of <i>cis</i> -MUFA replacement	38
Annex 8.	Results of carbohydrate replacement	39
Annex 9.	Risk of bias assessments	40
Annex 10.	GRADE evidence profiles	41
Reference	S	56

Acknowledgements

This systematic review was presented to, and reviewed by, the WHO Nutrition Guidance Expert Advisory Group (NUGAG) Subgroup on Diet and Health, which guided WHO in the interpretation of the results and in the development of the recommendations on intake of saturated fatty acids, taking into consideration not only those results, but also diverse values and preferences, balance of benefits and harms, resource implications, priority of the problem, equity and human rights, acceptability and feasibility.

Technical editing of the document was undertaken by Dr Hilary Cadman from Cadman Editing Services in Australia, and cover design and formatting of the document was undertaken by Ms Sue Hobbs from minimum graphics in New Zealand.

Abbreviations and acronyms

ApoA-I apolipoprotein A-I
ApoB apolipoprotein B
BMI body mass index

CHD coronary heart disease
CI confidence interval

cis-MUFA cis-monounsaturated fatty acids
cis-PUFA cis-polyunsaturated fatty acids

CVD cardiovascular disease

GRADE Grading of Recommendations Assessment, Development and Evaluation

HDL high-density lipoprotein

LDL low-density lipoprotein

NCDs noncommunicable diseases

NUGAG WHO Nutrition Guidance Expert Advisory Group

PICO population, intervention, comparator and outcome

RCT randomized controlled trial

SFA saturated fatty acids

TFA trans-fatty acids

WHO World Health Organization

1. Introduction

1.1 Background

Fats in the diet mainly consist of triglyceride, a molecule composed of three fatty acids and a glycerol backbone. Fatty acids differ in several aspects. First, they are characterized by the number of double bonds. Saturated fatty acids (SFA) have no double bonds, while monounsaturated fatty acids (MUFA) have one double bond and polyunsaturated fatty acids (PUFA) have two or more double bonds. These double bonds have either the cis or trans configuration. Most unsaturated fatty acids in the diet have the cis configuration, but trans-fatty acids (TFA) are also present. Second, the position of the double bond varies. Third, fatty acids differ in chain length, though the number of carbon atoms is usually an even number. The most abundant SFA in the diet have 16 (C16:0; palmitic acid) or 18 (C18:0; stearic acid) carbon atoms, while smaller proportions of SFA have 14 (C14:0; myristic acid) or 12 (C12:0; lauric acid) carbon atoms. Some fats (e.g. coconut oil and dairy fat) also contain fatty acids with fewer than 12 carbon atoms. The most abundant cis-MUFA is oleic acid (C18:1), and the most abundant cis-PUFA are linoleic acid (C18:2n-6) and α -linolenic acid (C18:3n-3). Existing evidence suggests that the intake of fatty acids is a major determinant of the serum lipid and lipoprotein profile.

1.2 Objectives

The aim of this systematic review was to assess the effect of modifying SFA intake on serum lipid and lipoprotein levels by exchanging SFA with *cis*-MUFA, *cis*-PUFA or carbohydrates, in order to inform and contribute to the development of updated WHO recommendations on SFA intake. Effects of TFA were not considered in this analysis.

2. Methods

This systematic review and regression analysis was conducted in accordance with the WHO guideline development process (1). As part of the evidence review, results of the regression analysis were evaluated using the methodology of the Grading of Recommendations Assessment, Development and Evaluation (GRADE) working group (2). Evidence summaries and GRADE assessments were discussed and reviewed by the WHO Nutrition Guidance Expert Advisory Group (NUGAG) Subgroup on Diet and Health, as part of the WHO guideline development process. The PICO (population, intervention, comparator and outcome) questions (Annex 1) and priority health outcomes (Annex 2) guiding this review were discussed and developed by the NUGAG Subgroup on Diet and Health.

This systematic review and analysis is an update of the results of an earlier published review and analysis (3).

2.1 Criteria for selecting studies to include in this review

2.1.1 Study characteristics

Study design

The review included only studies that were designed to eliminate the effect of nonspecific drifts of the outcome variables with time. Elimination of the effect could be achieved by feeding the different diets side-by-side (parallel design) or by giving the diets to the volunteers in random order (crossover or Latin square design). "Before-and-after" (sequential) designs do not eliminate this effect and were therefore excluded. Dietary periods had to be at least 13 days, because time is otherwise too short for serum lipids and lipoproteins to reach a new steady-state situation (4, 5).

Diets and interventions

Only studies with a thorough daily control of food intake were selected. Protein and alcohol intake had to be constant and fatty acids had to be exchanged for other fatty acids or for carbohydrates. Possible effects of protein and alcohol on the serum lipoprotein profile could therefore not be estimated. Other concomitant interventions (e.g. those targeting weight loss) were not allowed. Daily cholesterol intake between diets within a study had to be comparable (<100 mg difference). Diets that focused on (hydrogenated) very long chain (n-3) PUFA (fish oils) were excluded. Therefore, total PUFA in these studies can be considered to equal PUFA with 18 carbon atoms (linoleic acid plus α -linolenic acid). Studies focusing on medium-chain fatty acids (MCFA) or behenic acid were also excluded, because their number was too limited to allow proper statistical analyses. Only studies with a reported TFA intake of 2% of total energy intake or less were included. If TFA was not reported, it was assumed to be less than 2%. Estimates for the effects of the various fatty acids on serum lipids and lipoproteins were based on within-study comparisons (see **Section 2.2.4**), therefore studies that could only provide one data point based on the inclusion criteria were also excluded.

Participants

Only studies with apparently healthy adult subjects (aged > 17 years), who did not suffer from gross disturbances of lipid metabolism or from diabetes, were considered.

2.1.2 Outcomes

The outcomes assessed in this analysis were serum lipids and lipoproteins, including total cholesterol, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, triglyceride, LDL cholesterol to HDL cholesterol ratio, total cholesterol to HDL cholesterol ratio, triglyceride to HDL cholesterol ratio, apolipoprotein A-I (ApoA-I) and apolipoprotein B (ApoB) (Annex 2).

2.2 Data collection and analysis

2.2.1 Identification of studies

Search strategy

As indicated in **Section 2**, this analysis is an update of the results of an earlier published analysis (3). For the original analysis, controlled dietary trials published in English between January 1970 and December 1998 were identified through a computer-assisted literature search. Reference lists of identified papers were hand-searched for additional relevant citations. In total, 60 studies were identified that met the inclusion criteria. In 2009, a computer-assisted literature search was performed for articles published between January 1999 and December 2008, which brought the total data set to 83 studies. Finally, in January 2014, a computer-assisted literature search was performed in the PubMed database for articles published between January 2009 and December 2013. Search terms can be found in **Annex 3**. After scanning, an additional eight articles were identified, providing a total of 91 studies.

Selection of studies

A study was excluded if it was evident from the title or abstract that the study did not meet the inclusion criteria (e.g. the study addressed the effects of fish oils only, was not adequately controlled, was not an intervention study, etc.). Full texts of the remaining citations were reviewed for inclusion.

2.2.2 Data extraction and management

For studies meeting the inclusion criteria, data were extracted using standard data extraction forms. Data were then transferred in duplicate to Microsoft Excel. Typographical errors were corrected and the data were analysed for consistency (e.g. sum of fatty acids, sum of percentage of energy from the macronutrients, etc.). Each data point consisted of the fatty acid and carbohydrate composition of a particular diet and the mean serum lipid or lipoprotein concentration or ratio of a group of subjects, as obtained at the end of a dietary period. For parallel designs, serum parameters were adjusted for differences between the intervention groups at baseline.

2.2.3 Assessment of risk of bias in included studies

Risk of bias was assessed for each included study through identification and extraction of relevant information on study design and conduct. The following areas, discussed below, can lead to bias (6), and were included for assessment, each being assigned a *low*, *high* or *unclear* risk of bias:

- random sequence generation
- allocation concealment
- ▶ blinding of participants and personnel
- ▶ incomplete outcome data
- selective reporting
- other sources of bias.

Random sequence generation

For each included study, it was determined whether randomization was employed and if so, whether the method used to generate the randomization sequence was described in sufficient detail to allow an assessment of whether it would have produced comparable groups. Studies were categorized as one of the following in relation to risk of bias:

low – if a truly random process was used (e.g. random number table or computer random number generator); or a crossover study design was used, such that both groups received both the intervention and control treatment, and thus observed differences were unlikely to be a result of group differences;

- ► *high* if a non-random process was used (e.g. odd or even date of birth, or hospital or clinic record number), *or* randomization was not used; or
- ▶ *unclear* if the study did not specify whether randomization was used at all, or did not provide enough detail to determine whether the process was truly random.

Allocation concealment

For each included study, it was determined whether the method used to conceal the allocation sequence (in randomized studies) was described in sufficient detail so as to determine whether intervention allocation could have been foreseen in advance of or during recruitment, or changed after assignment. Studies were categorized as one of the following in relation to risk of bias:

- ▶ low if methods such as telephone or central randomization, consecutively numbered sealed opaque envelopes and so on were used; or if the studies had a crossover design or no randomization (in which case, allocation concealment is not relevant and thus does not present a source of bias);
- ► *high* if methods such as open allocation, unsealed or non-opaque envelopes, alternation, date of birth and so on were used; or
- unclear if the study did not specify whether allocation concealment was used at all, or did not provide enough detail to determine whether the process was sufficient to prevent knowledge of assignment.

Blinding of participants and personnel

For each included study, the methods used, if any, to blind study participants and personnel from knowledge of which intervention a participant received, were identified. Studies were judged to be at low risk of bias if they were blinded, or if it was deemed that the lack of blinding was unlikely to have affected the results. Studies were categorized as *low*, *high* or *unclear* risk of bias separately for:

- participants
- personnel
- ▶ outcome assessments.

Incomplete outcome data

For each included study, the completeness of data was determined, including attrition and exclusion of data from the analysis. It was further determined whether attrition and exclusions were reported, the numbers included in the analysis at each stage (compared with the total number of participants), reasons for attrition or exclusion (where reported), and whether missing data were balanced across groups or were related to outcomes. Studies were categorized as one of the following in relation to risk of bias:

- ▶ *low* if few drop-outs or losses to follow-up were noted or an intention-to-treat analysis was possible;
- high if there was significant loss to follow-up that was not addressed in terms of comparability across intervention and control groups, or data were not adjusted for missing data, or there were wide differences in exclusions between groups, whether or not intention-to-treat analysis was used; or
- ▶ *unclear* if losses to follow up or exclusions were not sufficiently reported to determine whether the process was sufficient.

Selective reporting

For each included study, an attempt was made to determine whether there was selective outcome reporting. Studies were categorized as one of the following in relation to risk of bias:

► *low* – if it was clear that all of the prespecified outcomes and all expected outcomes of interest to the review had been reported;

- high if not all prespecified outcomes had been reported; one or more reported primary outcomes had not been prespecified; outcomes of interest were reported incompletely and so could not be used; or results of a key outcome that would have been expected to have been reported were not reported; or
- ▶ *unclear* if the information given was insufficient to judge whether or not outcomes were selectively reported.

Other sources of bias

For each included study, other possible sources of bias were identified, such as potential differences in the groups at baseline, evidence of treatment compliance, residual confounding and other problems that could put it at risk of bias.

2.2.4 Analytical methods

Calculations and conversions

Plasma values for total and HDL cholesterol were multiplied by 1.030 and those for triglyceride by 1.029 to convert them to serum values (7). LDL cholesterol concentrations were calculated using the Friedewald equation (8). For the sake of uniformity, the total cholesterol to HDL cholesterol ratio, the LDL cholesterol to HDL cholesterol ratio, the triglyceride to HDL cholesterol ratio, and the LDL cholesterol concentration for all studies were recalculated, even if these values were reported by the study authors. Plasma values for ApoA-I and ApoB were also multiplied by 1.030 to convert them to serum values.

Dietary fat contains on average 96% by weight as fatty acids; the other 4% are glycerol and other lipids. For publications in which the intakes of the various fatty acid classes had been normalized so as to add up to 100% of total fat, the intakes were converted back into true fatty acid intakes by multiplying them by 0.96.

Statistical analysis

Multiple regression analyses were performed to predict the mean serum lipid or lipoprotein concentration or ratio (the dependent variable) of a group of subjects from the changes in percentage of total energy intake of fatty acids or carbohydrates in the diets (the independent variables). As indicated in **Section 2.1.1**, within each study fatty acids of the experimental diets were exchanged for other fatty acids or for carbohydrates.

In order to estimate the effects on serum lipids of both decreasing SFA intake (via replacement with *cis*-PUFA, *cis*-MUFA or carbohydrates) and increasing SFA intake (via replacement of *cis*-PUFA, *cis*-MUFA or carbohydrates with SFA), four models were generated. A fifth model estimated the effects of individual SFA. All five models used absolute lipid or lipoprotein concentrations or ratios on each diet as dependent variables. A dummy variable for each study was introduced into the model, to ensure that only withinstudy diet-induced differences were analysed. The estimate for that dummy variable can be envisaged as the mean estimated serum lipid or lipoprotein parameter ("the intrinsic level"), when the participants from that study consumed a standardized fat-free diet. It varies between studies, due to differences in study population (e.g. genetic makeup, age, and body mass index [BMI]), but also by other factors such as the fibre, protein or cholesterol content of the background diet, which was constant within studies, but differed between studies.

In the first model (model 1), SFA served as the reference nutrient and the effects of a mixture of *cis*-PUFA, *cis*-MUFA and carbohydrates, on the dependent variable were estimated. The regression coefficients represent the predicted change in the mean serum lipid or lipoprotein concentration or a ratio when SFA intake decreases by 1% of total energy intake and that of *cis*-PUFA, *cis*-MUFA or carbohydrates increases by the same amount. Diets in which the fatty acid composition of a particular class of fatty acids diverged markedly from that in normal mixed diets, were excluded (e.g. diets specifically enriched in lauric acid or stearic acid). Including these data points would have resulted in less reliable estimates of the effects of a normal mixture of SFA, because evidence indicates that individual SFA have different effects on the serum lipoprotein profile *(9, 10)*.

In the second model (model 2), cis-MUFA served as the reference nutrient and the effects of a mixture of cis-PUFA, carbohydrates and SFA on the dependent variable were estimated. The regression coefficients in this model now represent the predicted change in the mean serum lipid or lipoprotein concentration or a ratio when cis-MUFA intake decreases by 1% of total energy intake and that of cis-PUFA, carbohydrates or SFA increases by the same amount.

In the third model (model 3), cis-PUFA served as the reference nutrient and the effects of a mixture of cis-MUFA, carbohydrates and SFA on the dependent variable were estimated. The regression coefficients in this model now represent the predicted change in the mean serum lipid or lipoprotein concentration or a ratio when cis-PUFA intake decreases by 1% of total energy intake and that of cis-PUFA, carbohydrates or SFA increases by the same amount.

In the fourth model (model 4), carbohydrate served as the reference nutrient and the effects of a mixture of cis-PUFA, cis-MUFA and SFA on the dependent variable were estimated. The regression coefficients in this model now represent the predicted change in the mean serum lipid or lipoprotein concentration or a ratio when carbohydrate intake decreases by 1% of total energy intake and that of cis-PUFA, cis-MUFA or SFA increases by the same amount.

The fifth model (model 5) estimated the effects of individual SFA. The proportions of energy from lauric, myristic, palmitic and stearic acids were used as independent variables, together with the proportions of energy from *cis*-PUFA and *cis*-MUFA. Thus, as in the fourth model above, carbohydrates served as the point of reference. Individual SFA of less than 12 carbon atoms, were not reported in all studies and were therefore not included in the analysis.

The validity of the regression models was examined in several ways. First, normality of the residuals was checked. If the residual was not normally distributed, the most extreme value(s) were excluded. This approach did not change conclusions, but resulted in narrower confidence intervals. Second, the influence of each separate observation on the estimated regression coefficients was assessed using the Cook's distance. Observations with a Cook's distance >0.4 were excluded in the final analysis. Third, visual inspection of plots did not suggest a relationship between residuals and the independent variables. This suggests that the differences between observed and predicted values (i.e. the residuals) did not depend on the absolute level of intake of a particular (class of) fatty acid(s). Results of residuals analysis for SFA intake (model 1) are provided in **Annex 4**. Furthermore, the observed and predicted values for LDL cholesterol concentrations were in excellent agreement (**Annex 5**). Each data point was weighed for the number of participants. All statistical analyses were carried out using SPSS version 23.

Subgroup and sensitivity analyses

Subgroup analyses were performed to examine whether responses to the diets were influenced by baseline lipid and lipoprotein concentrations, gender (by comparing results of studies carried out with only men versus those carried out in men and women or women only) or publication date (by comparing results of studies published before 1993 and in 1993 or later, because it was at approximately that time that the detrimental effects of TFA on the serum lipid and lipoprotein profile became known). Subgroup analysis by type of carbohydrate included in study diets was also planned. Sensitivity analysis was performed in which studies that used liquid formula diets were excluded.

3. Results

3.1 Search results

The initial search for articles published between January 2009 and December 2013 returned 629 potentially eligible articles. After removing citations based on title or abstract, the full texts of 66 articles were assessed for inclusion, and eight were included. Together with the 83 articles from previous searches, a total of 91 dietary trials were identified. Seven of these studies could not be used for the final calculations, because they yielded only one data point (as the intake of TFA in the other diets exceeded 2% of total energy intake) and were therefore excluded, leaving 84 studies. The flow of records through screening, exclusion and inclusion of studies is shown in **Figure 1**.

3.2 Included studies

Characteristics of the 84 included studies are summarized in **Table 1**. The studies yielded 211 diet data points and included 2353 volunteers, of which 65% were men (n=1538) and 34% were women (n=801). For two studies with a total of 14 subjects, the number of men and women was not specified. Forty-six studies were carried out in men only, three studies in women only, and 35 studies in men and women. The diets were fed for 13 to 91 days. Seventy-three studies used a crossover design and 11 studies a parallel design. Forty-seven studies were from the United States of America (USA); eight from the Netherlands; six from Canada; five from Denmark; three from the United Kingdom of Great Britain and Northern Ireland; two each from Israel, Germany, Malaysia, Norway or Spain; and one each from Finland, Italy, New Zealand, Austria and Sweden. Seventeen diets from six studies consisted of liquid formula diets. Seventy-three trials reported the mean age of their participants, which varied between 21 and 72 years (mean 38 years). For 66 studies, BMI values were reported and ranged from 20.0 to 28.6 kg/m² (mean 24.2 kg/m²). For serum total cholesterol (63 studies), mean pre-study levels ranged from 3.7 to 6.7 mmol/L (mean 5.1 mmol/L); for LDL cholesterol (54 studies), from 2.3 to 4.8 mmol/L (mean 3.3 mmol/L); for HDL cholesterol (53 studies), from 0.9 to 1.8 mmol/L (mean 1.2 mmol/L); and for triglyceride (57 studies), from 0.7 to 2.2 mmol/L (mean 1.2 mmol/L).

3.3 Effects of interventions

3.3.1 Analysis of total SFA intake

3.3.1.1 Characteristics of included studies

The 74 trials used to examine the effects of reducing or increasing total SFA intake on serum lipids and lipoproteins yielded 177 total diet data points. The number of diet data points included in the calculations varied from 102 (41 studies) for ApoA-I to 177 (74 studies) for total cholesterol. Mean intake of total fat on these 177 diets was 34.0% of total energy intake (range 4.5–53.0%); of SFA, 9.8% of total energy intake (range 1.6–24.4%); of cis-MUFA, 13.6% of total energy intake (range 1.6–39.8%); and of cis-PUFA, 8.4% of total energy intake (range 0.4–28.8%).

3.3.1.2 Effects of reducing SFA intake by replacing SFA with other nutrients

These effects were generated using SFA as the reference nutrient (model 1) as described in **Section 2.2.4**.

Replacement of SFA with cis-PUFA

Results from multiple regression analysis (weighted for N) are summarized in **Table 2** and demonstrate that for each 1% of dietary energy¹ as SFA replaced with an equivalent amount of *cis*–PUFA, there was a

¹ As a percentage of total energy intake

significant decrease¹ in total cholesterol of 0.064 mmol/L (95% CI: -0.070, -0.058), in LDL cholesterol of 0.055 mmol/L (95% CI: -0.061, -0.050), in HDL cholesterol of 0.005 mmol/L (95% CI: -0.006, -0.003), in triglyceride of 0.010 mmol/L (95% CI: -0.014, -0.007), in the total cholesterol to HDL cholesterol ratio of 0.034 (95% CI: -0.040, -0.028), in the LDL cholesterol to HDL cholesterol ratio of 0.034 (95% CI: -0.009, -0.002; P = 0.004), in ApoA-I of 4.9 mg/dL (95% CI: -7.3, -2.5) and in ApoB of 10.2 mg/dL (95% CI: -12.4, -8.1).

Replacement of SFA with cis-MUFA

Results from multiple regression analysis (weighted for N) are summarized in **Table 2** and demonstrate that for each 1% of dietary energy as SFA replaced with an equivalent amount of *cis*-MUFA, there was a:

- ▶ significant decrease¹ in total cholesterol of 0.046 mmol/L (95% CI: -0.051, -0.040), in LDL cholesterol of 0.042 mmol/L (95% CI: -0.047, -0.037), in HDL cholesterol of 0.002 mmol/L (95% CI: -0.004, 0.000; P = 0.014), in triglyceride of 0.004 mmol/L (95% CI: -0.007, -0.001; P = 0.022), in the total cholesterol to HDL cholesterol ratio of 0.027 (95% CI: -0.033, -0.022), in the LDL cholesterol to HDL cholesterol ratio of 0.029 (95% CI: -0.034, -0.024) and in ApoB of 7.8 mg/dL (95% CI: -9.5, -6.0); and
- ▶ non-significant decrease in the triglyceride to HDL cholesterol ratio of 0.002 (95% CI: -0.005, 0.002; P = 0.342) and in ApoA-I of 1.8 mg/dL (95% CI: -3.7, 0.1; P = 0.064).

Replacement of SFA with carbohydrates

Results from multiple regression analysis (weighted for N) are summarized in **Table 2** and demonstrate that for each 1% of dietary energy as SFA replaced with an equivalent amount of carbohydrates, there was a:

- significant decrease¹ in total cholesterol of 0.041 mmol/L (95% CI: −0.047, −0.035), in LDL cholesterol of 0.033 mmol/L (95% CI: −0.039, −0.027), in HDL cholesterol of 0.010 mmol/L (95% CI: −0.012, −0.008), in the LDL cholesterol to HDL cholesterol ratio of 0.007 (95% CI: −0.013, −0.001; P = 0.017), in ApoA-I of 7.0 mg/dL (95% CI: −9.0, −5.1) and in ApoB of 3.6 mg/dL (95% CI: −5.4, −1.7 mg/dL);
- ▶ significant increase¹ in triglyceride of 0.011 mmol/L (95% CI: 0.007, 0.014) and the triglyceride to HDL cholesterol ratio of 0.014 (95% CI: 0.010, 0.018); and
- ▶ non-significant increase in the total cholesterol to HDL cholesterol ratio of 0.001 (95% CI: -0.006, 0.007; P = 0.842).

3.3.1.3 Effects of increasing SFA intake by replacing SFA with other nutrients

These effects were generated using *cis*-PUFA (model 2), *cis*-MUFA (model 3) or carbohydrates (model 4) as the reference nutrients as described in **Section 2.2.4**. (The full results for each analysis are provided in **Annexes 6–8**).

Replacement of cis-PUFA with SFA

Results from multiple regression analysis (weighted for N) are summarized in **Table 3** and demonstrate that for each 1% of dietary energy as cis-PUFA replaced with an equivalent amount of SFA, there was a significant increase¹ in total cholesterol of 0.066 mmol/L (95% CI: 0.060, 0.073), in LDL cholesterol of 0.058 mmol/L (95% CI: 0.052, 0.064), in HDL cholesterol of 0.005 mmol/L (95% CI: 0.004, 0.007), in triglyceride of 0.010 mmol/L (95% CI: 0.006, 0.014), in the total cholesterol to HDL cholesterol ratio of 0.034 (95% CI: 0.027, 0.041), in the LDL cholesterol to HDL cholesterol ratio of 0.035 (95% CI: 0.028, 0.041), in the triglyceride to HDL cholesterol ratio of 0.004 (95% CI: 0.001, 0.008; P = 0.026), in ApoA-I of 6.3 mg/dL (95% CI: 3.9, 8.7) and in ApoB of 10.3 mg/dL (95% CI: 7.7, 12.8).

Replacement of cis-MUFA with SFA

Results from multiple regression analysis (weighted for N) are summarised in **Table 3** and demonstrate that for each 1% of dietary energy as *cis*-MUFA replaced with an equivalent amount of SFA, there was a:

¹ P < 0.001 unless otherwise noted

- ▶ significant increase¹ in total cholesterol of 0.049 mmol/L (95% CI: 0.043, 0.055), in LDL cholesterol of 0.045 mmol/L (95% CI: 0.039, 0.051), in HDL cholesterol of 0.003 mmol/L (95% CI: 0.001, 0.004), in triglyceride of 0.004 mmol/L (95% CI: 0.000, 0.007; P = 0.041), in the total cholesterol to HDL cholesterol ratio of 0.028 (95% CI: 0.021, 0.034), in the LDL cholesterol to HDL cholesterol ratio of 0.030 (95% CI: 0.024, 0.036), in ApoA-I of 2.7 mg/dL (95% CI: 0.7, 4.8) and in ApoB of 8.1 mg/dL (95% CI: 6.1, 10.1); and
- ► non-significant increase in the triglyceride to HDL cholesterol ratio of 0.001 (95% CI: -0.003, 0.004; P = 0.680).

Replacement of carbohydrates with SFA

Results from multiple regression analysis (weighted for N) are summarized in **Table 3** and demonstrate that for each 1% of dietary energy as carbohydrates replaced with an equivalent amount of SFA, there was a:

- ▶ significant increase¹ in total cholesterol of 0.045 mmol/L (95% CI: 0.038, 0.051), in LDL cholesterol of 0.036 mmol/L (95% CI: 0.030, 0.043), in HDL cholesterol of 0.011 mmol/L (95% CI: 0.010, 0.013), in the LDL cholesterol to HDL cholesterol ratio of 0.007 (95% CI: 0.001, 0.014; *P* = 0.033), in ApoA-I of 8.4 mg/dL (95% CI: 6.4, 10.5) and in ApoB of 3.7 mg/dL (95% CI: 1.7, 5.8);
- ▶ significant decrease¹ in triglyceride of 0.012 mmol/L (95% CI: -0.015, -0.008) and in the triglyceride to HDL cholesterol ratio of 0.016 (95% CI: -0.020, -0.012); and
- ▶ non-significant decrease in the total cholesterol to HDL cholesterol ratio of 0.002 (95% CI: -0.009, 0.005; P = 0.553).

3.3.1.4 SFA intake at less than 10% of total energy intake

The population nutrient intake goal for SFA recommended by the joint WHO/FAO expert consultation (11) is less than 10% of total energy intake. One of the PICO questions which guided this systematic review was, therefore, designed to look at the effects of SFA consumption above and below 10% of total energy intake. Effects of reducing or increasing SFA intake on serum lipids and lipoproteins were observed across a wide range of SFA intakes, from 1.6 to 24.4% of total energy intake. Of the 177 data points used in the multiple regression, 113 included an SFA intake component of less than 10% of total energy intake and 65 included intakes of less than 8%. As noted in **Section 2.2.4**, analysis of the residuals of the regression line indicates that the relationship between a reduction or increase in SFA intake and effect on serum lipids and lipoproteins is linear with a consistent effect on serum lipids and lipoproteins across the entire range of SFA intakes. The results of the regression analysis therefore suggest reducing SFA intake to less than 10% of total energy intake may have additional benefit in terms of improving the overall serum lipoprotein profile when replacing SFA with *cis*-PUFA, *cis*-MUFA – and to a lesser extent carbohydrates – relative to higher intakes. Similarly, the results suggest a negative effect on the overall serum lipoprotein profile when increasing SFA intake from a starting point of less than 10% of total energy intake.

3.3.2 Subgroup and sensitivity analyses for total SFA

Results for subgroup and sensitivity analyses conducted with carbohydrates as the reference nutrient are described below. Results of subgroup and sensitivity analyses with other nutrients serving as the reference nutrient also did not show any significant differences between subgroups or when specified studies were removed for sensitivity analysis.

Subgroup analysis by type of carbohydrate was also planned. However, the number of studies reporting dietary data in sufficient detail to be able to determine with certainty the types of carbohydrates included in the study diets was limited, and therefore subgroup analysis could not be performed.

¹ P < 0.001 unless otherwise noted

3.3.2.1 Baseline levels

As described in **Section 2.2.4**, the estimate for the dummy variable in the regression model can be envisaged as the mean estimated serum lipid level when the participants from that study consumed a standardized fat-free diet. This estimate is a constant within studies, but differs between studies; that is, it can be considered a proxy for baseline lipid concentrations.

To examine whether baseline levels were related to responses, subgroup analyses were performed, in which the studies were split into low and high baseline groups based on the median level as estimated for each parameter based on model 4 in **Section 2.2.4**. The median levels when subjects consumed a standardized fat-free diet were as follows: total cholesterol 4.45 mmol/L, LDL cholesterol 2.89 mmol/L, HDL cholesterol 0.97 mmol/L, triglyceride 1.48 mmol/L, the total cholesterol to HDL cholesterol ratio 4.36, the LDL cholesterol to HDL cholesterol ratio 2.76 and the triglyceride to HDL cholesterol ratio 1.36. These analyses were not performed for ApoB and ApoA-I, because the number of studies in each subgroup was too small to provide reliable estimates for the regression coefficients.

Effect estimates

Effect estimates are presented in **Table 4**. The direction and statistical significance of the estimates did not depend on baseline levels. Effects, however, were in general more pronounced at higher baseline levels.

3.3.2.2 Gender

Thirty-eight studies were carried out in men only, 34 studies in men and women, and two studies in women only. These analyses were not performed for ApoB and ApoA-I, because the number of studies in each subgroup was too small to provide reliable estimates for the regression coefficients.

Effect estimates

Effect estimates are presented in **Table 5**. The direction and statistical significance of the estimates did not depend on gender. For total cholesterol, LDL cholesterol and HDL cholesterol, effects of a mixture of SFA in particular were less pronounced in studies that included men only.

3.3.2.3 Year of publication

In 1990, the detrimental effects of TFA on the serum lipoprotein profile were published for the first time. This may have resulted in an increasing awareness of the need to better analyse and report the intake of TFA of the study diets. Thirty-four studies were published before 1993 and 40 studies in 1993 or later. These analyses were not performed for ApoB and ApoA-I, because the number of studies in each subgroup was too small to provide reliable estimates for the regression coefficients.

Effect estimates

Effect estimates are presented in **Table 6**. The direction and statistical significance of the estimates did not depend on the year of publication. Also, the magnitude of the estimates was in good agreement, although effects of *cis*-PUFA on total cholesterol, LDL cholesterol and the LDL cholesterol to HDL cholesterol ratio were higher for studies published in 1993 or later.

3.3.2.4 Liquid formula diets

Eleven diets from five studies consisted of liquid formula diets. To examine the impact of these diets on the outcomes, analyses were repeated by excluding these studies.

Effect estimates

Effect estimates are presented in **Table 7** and do not suggest that removing studies that employed liquid formula diets substantially changed the results.

3.3.3 Analysis of individual SFA

These effects were generated using carbohydrates as the reference nutrient (model 5) as described in **Section 2.2.4**.

3.3.3.1 Characteristics of included studies

The 52 trials used to examine the effects of the individual SFA on serum lipids and lipoproteins yielded 134 diet data points. The number of diet data points included in the calculations varied from 88 for ApoA-I (34 studies) to 134 (51 studies) for total cholesterol. Mean intake of fat on these 134 diets was 35.6% of total energy intake (range 19.7–53.0%); of SFA, 12.0% of total energy intake (range 1.6–28.9%); of lauric acid (C12:0), 1.2% of total energy intake (range 0.0–16.9%); of myristic acid (C14:0), 1.2% of total energy intake (range 0.0–14.3%); of palmitic acid (C16:0), 5.9% of total energy intake (range 1.0–20.8%); and of stearic acid (C18:0), 2.8% of total energy intake (range 0.3–16.5%).

3.3.3.2 Effect estimates

Results from multiple regression analysis are summarized in **Table 8**. They demonstrate that for each 1% of dietary energy as carbohydrates replaced, there was a:

- significant increase¹ in:
 - ► total cholesterol when carbohydrates were replaced with lauric acid (0.029 mmol/L; 95% CI: 0.014, 0.045), myristic acid (0.060 mmol/L; 95% CI: 0.042, 0.077) or palmitic acid (0.041 mmol/L; 95% CI: 0.030);
 - ► LDL cholesterol when carbohydrates were replaced with lauric acid (0.017 mmol/L; 95% CI: 0.003, 0.031; *P* = 0.019), myristic acid (0.044 mmol/L; 95% CI: 0.028, 0.060) or palmitic acid (0.036 mmol/L; 95% CI: 0.026, 0.046);
 - ► HDL cholesterol when carbohydrates were replaced with lauric acid (0.019 mmol/L; 95% CI: 0.016, 0.023), myristic acid (0.021 mmol/L; 95% CI: 0.017, 0.025) or palmitic acid (0.010 mmol/L; 95% CI: 0.007, 0.013);
 - ► the LDL cholesterol to HDL cholesterol ratio when carbohydrates were replaced with palmitic acid (0.013; 95% CI: 0.005, 0.021; P = 0.002); and
- ► significant decrease¹ in:
 - ▶ triglyceride when carbohydrates were replaced with lauric acid (-0.015 mmol/L; 95% CI: -0.023, -0.007), myristic acid (-0.011 mmol/L; 95% CI: -0.020, -0.002; P = 0.018) or palmitic acid (-0.011 mmol/L; 95% CI: -0.017, -0.006);
 - ► the total cholesterol to HDL cholesterol ratio (-0.035; 95% CI: -0.048, -0.022) and the LDL cholesterol to HDL cholesterol ratio (-0.024; 95% CI: -0.036, -0.013) when carbohydrates were replaced with lauric acid; and
 - ► the triglyceride to HDL cholesterol ratio when carbohydrates were replaced with lauric acid (-0.024; 95% CI: -0.032, -0.017), myristic acid (-0.018; 95% CI: -0.027, -0.010) or palmitic acid (-0.015; 95% CI: -0.020, -0.009).

No significant associations were observed in:

- ► the total cholesterol to HDL cholesterol ratio when carbohydrates were replaced with myristic or palmitic acid;
- ► the LDL cholesterol to HDL cholesterol ratio when carbohydrates were replaced with myristic acid; or
- any serum lipid or ratio when carbohydrates were replaced with stearic acid.

¹ P < 0.001 unless otherwise noted

3.3.4 Subgroup and sensitivity analyses for individual SFA

3.3.4.1 Baseline levels

These analyses were not performed for individual SFA, because the number of studies in each subgroup was too small to provide reliable estimates for the regression coefficients.

3.3.4.2 Liquid formula diets

Twelve diets from four studies consisted of liquid formula diets. To examine the impact of these diets on the outcomes, analyses were repeated by excluding these studies. None of the diets that used liquid formula diets reported ApoB and ApoA-I concentrations; thus these analyses were not performed.

Effect estimates

Effect estimates are presented in **Table 9** and do not suggest that removing studies that employed liquid formula diets substantially changed the results.

3.3.4.3 Gender

These analyses were not performed for individual SFA, because the number of studies in each subgroup was too small to provide reliable estimates for the regression coefficients.

3.3.4.4 Year of publication

These analyses were not performed for individual SFA, because the number of studies in each subgroup was too small to provide reliable estimates for the regression coefficients.

3.4 Quality of the evidence

Some of the trials with parallel design were assessed as having unclear risk of bias in terms of randomization because the randomization procedure was not described adequately. Trials with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were randomized, as all participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have a significant, systematic effect on study results. Blinding was also not deemed to be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some trials may have been able to distinguish between intervention and control diets, this was not expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were determined to have a low risk of bias. Bias assessments for each study can be found in **Annex 9**.

The assessment of the quality of evidence for priority outcomes is found in the GRADE evidence profiles (**Annex10**). The quality of evidence for an effect of replacing SFA with *cis*-PUFA, *cis*-MUFA, or carbohydrates on all outcomes was judged to be high, except for ApoA-I and the triglyceride to HDL cholesterol ratio when replacing SFA with *cis*-MUFA and the total cholesterol to HDL cholesterol ratio when replacing SFA with carbohydrates, which were both judged as moderate due to serious imprecision. The quality of evidence for an effect of replacing *cis*-PUFA, *cis*-MUFA or carbohydrates with SFA on all outcomes was judged to be high, except for the triglyceride to HDL cholesterol ratio when replacing *cis*-MUFA with SFA, and the total cholesterol to HDL cholesterol ratio when replacing carbohydrates with SFA, which was judged as moderate due to serious imprecision. The quality of evidence was not assessed for outcomes in the analyses of individual SFA.

4. Strengths and limitations of review

4.1 Strengths

A strength of this study was that a large number of strictly-controlled dietary trials were identified and included in the regression analysis. Most of these studies were of relatively short duration (3–5 weeks), although long enough for serum lipid and lipoproteins concentrations to reach a new steady state. Tight control of dietary intake during the relatively short study period minimized non-compliance and other issues that often affect longer-term behaviour-change studies, increasing confidence in the results of the regression analysis.

Another strength is that the included studies cover a wide range of SFA intakes, from 1.6–24.4% of total energy intake, which increased the likelihood of detecting robust effects. Results are consistent across the entire range of intakes, suggesting that they could apply to a variety of populations with different SFA intakes.

The use of multiple regression allowed for assessment of the differential effects of replacing SFA with various nutrients, rather than simply estimating the effects of reducing or increasing SFA intake without regard to the nature of replacement. This is important, as a number of studies, including the original analysis on which this one is based, have shown that the effect of SFA reduction on serum lipids and lipoproteins is highly dependent on the nature of replacement.

4.2 Limitations

Inclusion of only those studies with strictly controlled diets greatly reduced the chance that dietary factors other than those being studied contributed to the changes observed in serum lipids and lipoproteins. This approach, while valuable in assessing specific effects of modifying SFA intake through exchange of specific nutrients, does not provide a clear picture of what might happen in real world settings in which modification of SFA intake might be accompanied by other changes in diet.

As indicated in **Section 4.1**, and described in the results (**Section 3**), different replacement scenarios have different effects on serum lipids and lipoproteins. Carbohydrate replacements as assessed in this analysis, included a mixture of monosaccharides, disaccharides and polysaccharides. Subgroup analysis by type of carbohydrate could not be performed because of the limited number of number of studies providing sufficient dietary information to determine, exactly, the composition of the carbohydrates in the diets. Assessment of the effects on the serum lipoprotein profile of replacing SFA with different types of carbohydrates would have been informative, given that a previous meta-analysis has suggested that diets with a low glycaemic index reduced total cholesterol and LDL cholesterol as compared with diets with a high glycaemic index (*12*). Furthermore, in an analysis of two large cohort studies, isocaloric replacement of SFA by carbohydrates from added sugars or refined carbohydrates was not associated with a change in risk of coronary heart disease, whereas replacement with carbohydrates from whole grains was related to a lower risk (*13*).

Lastly, although the analysis of individual SFA provided results for four common SFA in the diet, the intakes of lauric and myristic acid in included studies was generally quite low (i.e. mean of 1.2% of total energy intake). Thus, to obtain more insight into the effects of these two SFA on the serum lipoprotein profile at higher intakes, more well-controlled intervention studies at higher intakes are needed. Also, effects of SFA with less than 12 carbon atoms or more than 18 carbon atoms could not be estimated due to lack of information.

5. Conclusion

Results of the multiple regression analysis indicated that effects on the serum lipoprotein profile of reducing SFA intake by replacing a mixture of SFA with cis-PUFA (predominantly linoleic acid and α -linolenic acid) or cis-MUFA (predominantly oleic acid) were more favourable than replacing SFA with a mixture of carbohydrates. For total and LDL cholesterol and triglycerides in particular, the most favourable effects were observed for cis-PUFA. These results are consistent across a wide range of SFA intakes including intakes of less than 10% of total energy intake.

Differences in effects of the individual SFA on the serum lipoprotein profile were observed. Compared with a mixture of carbohydrates, an increased intake of lauric, myristic or palmitic acid raised serum total, LDL and HDL cholesterol levels, and lowered triglyceride levels, while increased intake of stearic acid did not appear to have a significant effect on these or other serum lipid values. Lauric acid alone reduced the total cholesterol to HDL cholesterol and LDL cholesterol to HDL cholesterol ratios as compared with a mixture of carbohydrates.

No significant gender-specific differences were observed regarding SFA intake and effects on serum lipids and lipoproteins, nor were the observed results systematically affected by dates of study publication, or inclusion of liquid diets in studies. In addition, conclusions did not change if subjects were stratified for baseline levels at the start of the study. It was not possible to perform subgroup analysis by type of carbohydrate.

15

6. Tables

Table 1. Characteristics of included studies

Reference and	Reference and		Co	mposition	1 ¹				
country	Study design	Diet	s	М	Р	Т	Participants		Funding
Mensink 1987 (14) ² Mensink 1989 (15) The Netherlands	Randomized parallel design with two interventions Experimental period: 35 days	1. 2.	6.7 9.8	9.3 24.0	5.2 5.1		Initial: 57, final: 48 Reason for loss: influenza (n=3), change in smoking habits (n=2), weight loss (n=4)	Diet 1: 12 men, 12 women Diet 2: 12 men, 12 women Mean age: 27 years	The Commission of the European Communities
Mattson 1985 <i>(16)</i> USA	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3.	19.1 3.3 4.3	15.4 28.2 5.6	3.9 6.9 28.1		Initial: 12, final: 12 No dropouts reported	• 12 men • Mean age: 59 years	Veterans Administration National Institutes of Health Moss Heart Foundation
Grundy 1986 A (17) USA	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	3.8 6.4	26.9 6.4	7.7 6.4		Initial: 7, final: 7 No dropouts reported	Sex not reported Mean age: 58 years	Veterans Administration National Institutes of Health Southwestern Medical Foundation Mead Johnson and Company Moss Heart Foundation
Brussaard 1980 <i>(18)</i> The Netherlands	Randomized parallel design with four interventions Experimental period: 35 days	1. 2. 3. 4.	8.0 10.0 11.0 18.0	10.0 8.0 8.0 16.0	3.0 11.0 19.0 3.0		Initial: 60, final: 60 No dropouts reported	 37 men and 23 women Diet 1: 16 subjects Diet 2: 15 subjects Diet 3: 15 subjects Diet 4: 14 subjects Sex distribution not reported. Age: 18-28 years 	The Netherlands Heart Foundation
Brussaard 1982 <i>(19)</i> The Netherlands	Randomized parallel design with two interventions Experimental period: 91 days	1. 2.	9.0 7.0	10.0 8.0	11.0 4.0		Initial: 35, final: 35 No dropouts reported	Diet 1: 11 men and 6 women Diet 2: 12 men and 6 women Age: 19-30 years	The Netherlands Heart Foundation
Mensink 1989 (20) The Netherlands	Randomized parallel design with two interventions Experimental period: 35 days	1. 2.	12.9 12.6	15.1 10.8	7.9 12.7		Initial: 60, final: 58 No reason for loss reported	Diet 1: 14 men and 15 women Diet 2: 13 men and 16 women Mean age: 25 years	Netherlands Nutrition Foundation The Netherlands Heart Foundation The Netherlands Ministry of Health
Harris 1983 <i>(21)</i> USA	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	14.4 6.4	16.4 10.8	7.2 21.6		Initial: 7, final: 7 No dropouts reported	Sex not reported Mean age: 40 years	National Heart, Lung, and Blood Institute Clinical Research Center Grant
Becker 1983 (22) USA	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3.	2.7 4.0 20.3	29.2 15.1 13.7	6.5 17.5 4.1		Initial: 12, final: 12 No dropouts reported	12 men Mean age: 32 years	Clinical Research Center Program National Institutes of Health Corn Products

١	۰	-	١
Ġ			
ı	u	,	,

Reference and	Charles de stem		Co	mpositio	1 ¹		Dt.		For the s
country	Study design	Diet	s	М	Р	T	Partic	cipants	Funding
Bonanome 1988 <i>(23)</i> USA	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. 3.	19.6 19.9 3.1	14.9 15.2 30.6	3.7 3.2 4.7		Initial: 11, final: 11 No dropouts reported	• 11 men • Mean age: 72 years	Not reported
Grundy 1986 B (24) USA	Randomized crossover design with two interventions Experimental period: 60 days	1. 2.	9.6 9.6	12.5 9.6	16.3 9.6		Initial: 9, final: 9 No dropouts reported	9 men Mean age: 63 years	Veterans Administration / National Institutes of Health Southwestern Medical Foundation Moss Heart Foundation
Katan 1988 <i>(25)</i> The Netherlands	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	23.4 11.6	14.1 11.7	5.2 20.9	1.9 0.5	Initial: 54, final: 47 Reason for loss: illness, weight loss, poor compliance	24 men and 23 women Mean age: 44 years	The Netherlands Heart Foundation
Grande 1972 (26) USA	Randomized crossover design with four interventions Experimental period: 28 days	1. 2. 3. 4.	2.3 3.3 5.2 8.7	1.6 6.5 16.9 7.1	0.6 2.7 6.7 13.3		Initial: 48, final: 38 Reason for loss: transport to another institution, illness, poor eating habits	38 men Mean age: 56 years	Public Health Service Research Grants
Anderson 1976 A (27) ³ USA	Randomized crossover design with two interventions Experimental period: 14 days	1. 2.	19.6 4.8	8.4 5.1	5.2 22.7		Initial: 12, final: 12 No dropouts reported	• 12 men • Mean age: 21 years	Public Health Service Research Grants
Anderson 1976 B (27) ³ USA	Randomized crossover design with two interventions Experimental period: 14 days	1. 2.	19.4 4.8	8.4 5.1	5.1 22.9		Initial: 12, final: 12 No dropouts reported	• 12 men • Mean age: 21 years	Public Health Service Research Grants
Wolf 1983 <i>(28)</i> USA	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3.	19.2 9.7 14.4	9.5 9.6 7.2	9.7 9.5 7.2		Initial: 6, final: 6 No dropouts reported	6 men Mean age: 54 years	Not reported
Grundy 1988 <i>(29)</i> USA	Randomized crossover design with two interventions Experimental period: 42 days	1. 2.	6.7 6.7	25.9 6.7	5.8 5.8		Initial: 10, final: 10 No dropouts reported	• 10 men • Mean age: 64 years	Veterans Administration National Institutes of Health Moss Heart Foundation
Reiser 1985 (30) USA	Randomized crossover design with three interventions Experimental period: 35 days	1. 2. 3.	9.4 18.8 1.6	10.4 1.1 2.2	0.4 0.3 16.2		Initial: 19, final: 19 No dropouts reported	• 19 men • Mean age: 26 years	National Heart and Blood Vessel Research National Heart, Lung, and Blood Institute National Institutes of Health Clinical Research USDHS Grant Lipid Research Clinics National Live Stock and Meat Board The Texas Cattle Feeders Association The Standard Meat Co of Fort Worth
Laine 1982 <i>(31)</i> USA	Randomized crossover design with four interventions Experimental period: 20 days	1. 2. 3. 4. ⁴	8.6 2.6 3.0 2.4	7.7 4.6 4.2 6.1	1.8 11.1 11.1 7.3	2.8	Initial: 24, final: 24 No dropouts reported	• 13 men and 11 women • Mean age: 25 years	American Soy Bean Association General Clinical Research Centers Program National Institutes of Health

Kay 1985 (33) United Kingdom	with three interventions Experimental period: 35 days	2.	9.4 13.4	9.2 13.2	7.3 11.7		No dropouts reported	Mean age: 45 years	Notreported
McDonald 1989 <i>(34)</i> Canada	Randomized crossover design with two interventions Experimental period: 18 days	1. 2.	5.1 6.8	20.2 7.4	10.3 21.6		Initial: 8, final: 8No dropouts reported	• 8 men • Age: 19–32 years	Canola Council of Canada
Mensink 1990 <i>(35)</i> The Netherlands	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. ⁴ 3.	9.3 9.3 19.4	23.7 13.0 13.6	4.4 4.5 3.0	0.0 10.9 0.7	 Initial: 59, final: 59 No dropouts reported 	25 men and 34 women Mean age: 26 years	 The Netherlands Nutrition Foundation The Netherlands Ministry of Welfare, Public Health, and Cultural Affairs The Commission of the European Communities
Valsta 1992 <i>(36)</i> Finland	Randomized crossover design with two interventions Experimental period: 25 days	1. 2.	12.4 12.7	16.2 10.2	7.6 13.3		 Initial: 59, final: 59 No dropouts reported 	29 men and 30 womenMean age: 30 years	 Food Research Foundation The Ministry of Agriculture and Forestry The Yrjö Jahnsson Foundation The Academy of Finland The Finnish Cultural Foundation
Wahrburg 1992 (37) Germany	Randomized crossover design with two interventions Experimental period: 23 days	1. 2.	10.2 10.1	16.0 9.9	4.1 10.3		 Initial: 40, final: 38 Reason for loss: illness (n=1), genetic anomaly of lipid metabolism (n=1) 	• 21 men and 17 women • Mean age: 24 years	The Commission of the European Communities
Zock 1992 (38) The Netherlands	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. 3. ⁴	11.0 20.1 10.3	15.7 16.3 15.6	12.5 4.3 4.2	0.1 0.3 7.7	 Initial: 59, final: 56 Reason for loss: personal (n=1), illness (n=1), pregnancy (n=1) 	• 26 men and 30 women • Mean age: 24 years	Not reported
Wardlaw 1990 <i>(39)</i> ⁶ Kwon 1991 <i>(40)</i> USA	Randomized crossover design with two interventions Experimental period: 35 days	1. 2.	6.7 7.7	26.9 13.4	5.8 18.2		Initial: 22, final: 20Reason for loss: not reported	• 20 men • Mean age: 35 years	SVO Enterprises
Ginsberg 1990 <i>(41)</i> USA	Randomized parallel design with two interventions Experimental period: 70 days	1. 2.	9.0 8.8	10.6 17.2	10.0 10.1		 Initial: 39, final: 36 Reason for loss: allergy (n=1), poor compliance (n=2) 	Diet 1: 12 men Diet 2: 12 men Diet 3: 12 men Mean age: 23 years	The National Institutes of Heath Best Foods Kraft Inc. Bertolli
Chan 1991 <i>(42)</i> Canada	Randomized crossover design with four interventions Experimental period: 18 days	1. 2. 3. 4.	6.5 5.3 7.1 6.4	18.7 18.3 8.4 9.9	7.4 8.5 16.8 16.1		Initial: 8, final: 8 One subject dropped out and was replaced	8 men Age: 20-34 years	Canola Council of Canada Flax Council of Canada
Wardlaw 1991 <i>(43)</i> USA	Randomized parallel design with two interventions Experimental period: 56 days	1. 2.	6.7 6.7	21.1 8.6	10.6 21.1		 Initial: 34, final: 32 Reason for loss: medication (n=1), unusual lipid values (n=1) 	• Diet 1: 16 men • Diet 2: 16 men • Mean age: 33 years	The Procter & Gamble Company
Berry 1991 <i>(44)</i> Israel	Randomized crossover design with two interventions Experimental period: 84 days	1. 2.	8.0 7.1	15.9 6.2	7.5 16.0		 Initial: 26, final: 18 Reason for loss: drop out (n=4), incomplete blood sampling (n=4) 	18 men Mean age: not reported	The National Institutes of Health

• Initial: 12, final: 12

• 12 men

Not reported

9.6 9.2 7.2

Lewis 1981 (32)⁵

Randomized crossover design 1.

ľ				•	
	٦		٩	١	
۰	•	۰	•		

Reference and			Co	mpositio	1 ¹				
country	Study design	Diet	s	М	Р	Т	Partic	ipants	Funding
Berry 1992 <i>(45)</i> Israel	Randomized crossover design with two interventions Experimental period: 84 days	1. 2.	6.6 4.7	16.6 6.8	7.5 5.7		Initial: 26, final: 17 Reason for loss: not reported	• 17 men • Age: 18–24 years	The National Institutes of Health, Public Health Service
Kris-Etherton 1993 A (46) ⁷ USA	Randomized crossover design with four interventions Experimental period: 26 days	1. 2. 3. 4.	6.0 20.9 6.3 21.0	27.2 13.2 10.1 10.1	2.3 2.1 17.8 1.7		Initial: 19, final: 18 Reason for loss: not reported	• 18 men • Mean age: 26 years	The American Cocoa Research Institute The Pennsylvania Agricultural Experimental Station
Kris-Etherton 1993 B (46) ⁷ USA	Randomized crossover design with four interventions Experimental period: 26 days	1. 2. 3. 4.	20.7 21.0 20.5 20.0	12.1 13.3 12.3 10.4	1.8 2.1 1.7 1.6		Initial: 18, final: 15 Reason for loss: not reported	• 15 men • Mean age: 27 years	The American Cocoa Research Institute The Pennsylvania Agricultural Experimental Station
Tholstrup 1994 (47) Denmark	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	16.8 17.5	14.1 14.6	3.6 3.8		Initial: 59, final: 59 No dropouts reported	• 12 men • Mean age: 24 years	Danish Agricultural Ministry Danish Agricultural and Veterinary Research Council Danish Technical Research Council
Denke 1992 <i>(48)</i> USA	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. 3.	2.6 18.7 18.9	29.1 17.0 15.4	6.0 2.4 3.8		Initial: 14, final: 14No dropouts reported	• 14 men • Mean age: 63 years	Southwestern Medical Foundation Moss heart Foundation Veterans' Affairs National Heart, Lung, and Blood Institute
Bonanome 1992 (49) Italy	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	9.6 9.6	28.8 4.8	4.8 28.8		Initial: 11, final: 11 No dropouts reported	• 11 men • Mean age: 22 years	The European Economic Community
Judd 1994 (50) USA	Randomized crossover design with four interventions Experimental period: 42 days	1. 2. ⁴ 3. ⁴ 4.	14.0 13.6 13.1 20.1	16.4 14.6 13.5 10.9	5.9 5.7 6.4 5.8	0.7 3.7 6.4 0.7	Initial: 64, final: 58 Reason for loss: illness (n=1), no reason reported (n=1), other commitments (n=3), non-compliance (n-1)	29 men, 29 women Mean age: 43 years	Institute of Shortening and Edible Oils and its member companies
Sundram 1994 <i>(51)</i> Malaysia	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	15.0 14.1	11.0 11.6	3.2 3.5		Initial: 18, final: 17 Reason for loss: illness of family member (n=1)	• 17 men • Mean age: 21 years	Not reported
Tholstrup 1994 (52) Denmark	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. 3.	15.7 18.1 20.2	15.7 14.3 14.4	2.7 4.3 2.3		Initial: 15, final: 15 No dropouts reported	• 15 men • Mean age: 25 years	Danish Agricultural and Veterinary Research Council Danish Technical Research Council
Zock 1994 <i>(53)</i> The Netherlands	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. 3.	21.3 21.0 10.8	11.2 11.9 21.3	4.1 4.7 4.4	0.8 0.2 0.3	Initial: 59, final: 59 No dropouts reported	23 men and 36 women Mean age: 29 years	Foundation for Nutrition and Health Sciences
Barr 1992 <i>(54)</i> USA	Randomized parallel design with two interventions Experimental period: 49 days	1. 2.	9.0 12.2	13.2 10.8	7.8 6.5		Initial: 51, final: 48 Reason for loss: illness (n=1), poor compliance (n=2) 17 men received a diet that was not included in the meta-analysis	Diet 1: 15 men Diet 2: 16 men Mean age: 25 years	National Institutes of Health Best Foods, Kraft Inc. Bertolli

	_	١
5	_	
٠	S	,

Ginsberg 1994 (55) USA	Randomized parallel design with two interventions Experimental period: 42 days	1. 2.	8.9 9.1	8.4 13.2	11.4 6.4		Initial: 30, final: 30 No dropouts reported 12 men received a diet that was not included in the meta-analysis	• Diet 1:9 men • Diet 2:9 men • Mean age: 25 years	National Institutes of Health Best Foods, Kraft Inc. Bertolli
Judd 1988 (56)8 Marshall 1986 (57) USA	Randomized crossover design with two interventions Experimental period: 42 days	1. 2.	6.7 10.6	11.4 10.4	6.5 3.3		Initial: 24, final: 23 Reason for loss: personal	• 23 men • Age: 35–60 years	Not reported
Sundram 1995 (58) Malaysia	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	6.0 13.0	17.5 14.3	7.7 4.1		Initial: 24, final: 23 Reason for loss: not reported	23 men Mean age: 22 years	Not reported
lacono 1991 <i>(59)</i> USA	Randomized crossover design with two interventions Experimental period: 40 days	1. 2.	9.5 8.6	9.4 8.7	3.8 10.8		Initial: 11, final: 11 No dropouts reported	• 11 men • Mean age: 54 years	Not reported
Lichtenstein 1993 (60) ⁹ Lichtenstein 1994 (61) Lichtenstein 1994 (62) USA	Randomized crossover design with five interventions Experimental period: 32 days	1. 2. 3. 4. 5.	5.4 6.9 6.9 12.1 7.4	14.5 9.0 17.0 11.3 10.8	6.7 11.2 3.9 3.4 8.8		Initial: 15, final: 14 Reason for loss: scheduling problems (n=1)	6 men and 8 women Mean age: 63 years	US Department of Agriculture National Institutes of Health Uncle Bens, Inc
Dougherty 1995 (63) USA	Randomized crossover design with two interventions Experimental period: 40 days	1. 2.	10.8 9.1	9.5 9.5	7.9 7.0	0.6 1.0	Initial: 10, final: 10 No dropouts reported	• 10 men • Mean age: 37 years	Not reported
Marckmann 1992 <i>(64)</i> Denmark	Randomized crossover design with two interventions Experimental period: 14 days	1. 2.	15.4 13.5	11.8 8.2	6.0 4.7		Initial: 13, final: 13 No dropouts reported	6 men and 17 women Mean age: 26 years	The Danish Heart Foundation The Danish Health Insurance Foundation The Danish Agricultural and Veterinary Research Council
Howard 1995 <i>(65)</i> USA	Randomized crossover design with four interventions Experimental period: 42 days	1. 2. 3. 4.	8.2 8.0 9.4 9.5	14.2 12.1 8.5 5.7	3.1 4.8 7.2 12.5		Initial: 77, final: 63 Reason for loss: employment obligations (n=4), poor compliance (n=9), loss of blood samples (n=1)	30 men and 33 women Mean age: 46 years	National Heart, Lung, and Blood Institute Best Foods
Fielding 1995 A (66) ¹⁰ USA	Randomized parallel design with two interventions Experimental period: 28 days	1. 2.	10.3 15.3	16.5 15.4	8.5 5.8		Initial: 48, final: 42 Reason for loss: not reported (n=5), incomplete data (n=1)	42 menDiet 1: 21 menDiet 2: 21 menMean age: 29 years	National Institutes of Health Arteriosclerosis SCOR National Dairy Promotion and Research Board
Fielding 1995 B <i>(66)</i> ¹⁰ USA	Randomized parallel design with two interventions Experimental period: 28 days	1. 2.	10.0 16.7	14.9 12.7	9.9 4.7		Initial: 48, final: 42 Reason for loss: not reported (n=5), incomplete data (n=1)	42 menDiet 1: 20 menDiet 2: 22 menMean age: 29 years	National Institutes of Health Arteriosclerosis SCOR National Dairy Promotion and Research Board
Park 1996 (67) USA	Randomized crossover design with three interventions Experimental period: 28 days	1. ⁴ 2. ⁴ 3. ⁴	15.2 13.7 13.5	17.0 18.1 16.8	6.9 6.6 8.2		Initial: 18, final: 17 Reason for loss: not reported	• 17 men • Mean age: 26 years	The National Live Stock and Meat Board The Ohio Agricultural Experimental Station
Cater 1997 (68) USA	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	23.3 5.7	18.4 39.8	6.0 4.9		Initial: 9, final: 9 No dropouts reported	9 men Mean age: 66 years	NIH Endocrinology and Metabolism Training Grant NIH-NHLBI Clinical Investigator Award National Institutes of Health

ľ	١.	J

Reference and	Study design		Co	mpositio	1 ¹		Participants		Funding
country	Study design	Diet	s	М	Р	Т	Partic	ipants	Funding
Tholstrup 1998 (69) Denmark	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	19.1 24.4	11.6 7.7	4.5 5.2	1.6 0.1	Initial: 18, final: 18 No dropouts reported	• 18 men • Mean age: 25 years	The Danish Dairy Research Foundation The Danish Research Development Program for Food Technology
Mazier 1997 (70) Canada	Randomized crossover design with two interventions Experimental period: 13 days	1. 2.	11.0 10.9	24.0 9.2	4.1 17.9		Initial: 9, final: 9 No dropouts reported	• 9 men • Mean age: 26 years	The Heart and Stroke Foundation of British Columbia and Yukon
Ginsberg 1998 <i>(71)</i> USA	Randomized crossover design with three interventions Experimental period: 56 days	1. 2. 3.	14.4 8.6 5.8	12.5 12.5 12.5	5.8 5.8 5.8		Initial: 118, final: 103 Reason for loss: not reported	46 men and 57 women Mean age: 38 years	National Heart, Lung, and Blood Institute National Center for Research Resources
Müller 1998 <i>(72)</i> Norway	Randomized crossover design with three interventions Experimental period: 17 days	1. 2. ⁴ 3.	12.5 6.3 7.3	11.4 10.3 11.4	5.5 5.3 9.8	0.1 6.7 0.2	Initial: 30, final: 27 Reason for loss: not reported (n=2), poor compliance (n=1)	• 27 women • Mean age: 27 years	The Nordic Industrial Fund Mills DA
Hunter 2000 (72) United Kingdom	Randomized crossover design with three interventions Experimental period: 14 days	1. 2. 3.	17.6 6.8 7.3	13.9 25.0 14.4	4.5 4.5 14.4		Initial: 9, final: 6 Reason for loss: not reported	6 men Mean age: 28 years	Ministry of Agriculture, Food and Fisheries Scottish Executive Rural Affairs Department
Judd 2002 (74) ¹¹ Baer 2004 (75) USA	Randomized crossover design with six interventions Experimental period: 35 days	1. 2. 3. ⁴ 4. ⁴ 5. 6.	12.8 12.6 12.8 16.9 20.9 20.8	10.5 17.6 10.6 10.6 10.5 10.5	3.8 3.8 4.0 4.3 4.4 4.2	0.2 0.1 8.3 4.2 0.3 0.2	Initial: 54, final: 50 Reason for loss: not reported (n=3), poor compliance (n=1)	50 men Mean age: 42 years	Technical Committee on Dietary Lipids, International Life Sciences Institute
Vega-López 2006 (76)	Randomized crossover design with two interventions Experimental period: 35 days	1. 2.	14.8 6.4	10.9 15.4	3.5 8.7	0.6 1.0	Initial: 15, final: 15 No dropouts reported	• 5 men and 10 women • Mean age: 64 years	National Institutes of Health / US Department of Agriculture
Lichtenstein 1999 (77) USA	Randomized crossover design with five interventions Experimental period: 35 days	1. 2. 3. ⁴ 4. ⁴ 5. ⁴	7.3 8.6 8.4 8.6 8.5	8.1 8.0 9.9 8.5	12.5 13.5 11.1 8.1 6.3	0.6 0.9 3.3 4.2 6.7	Initial: 36, final: 36 No dropouts reported	18 men and 18 women Mean age: 63 years	National Institutes of Health US Department of Agriculture
Lovejoy 2002 <i>(78)</i> USA	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3. ⁴	5.9 10.9 7.2	14.7 8.8 7.6	6.3 6.4 4.0	0.0 0.0 7.0	Initial: 31, final: 25 Reason for loss: not reported	• 12 men and 13 women • Mean age: 28 years	US Department of Agriculture
Berglund 2007 <i>(79)</i> USA	Randomized crossover design with three interventions Experimental period: 49 days	1. 2. 3.	15.0 8.4 7.7	13.8 20.0 14.9	5.6 6.0 5.3		Initial: 110, final: 85 Reason for loss: not reported	• 52 men and 33 women • Mean age: 36 years	National Institutes of Health
Binkoski 2005 (80) USA	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3.	10.8 8.0 7.6	14.3 16.5 13.6	7.5 4.1 7.4		Initial: 31, final: 31 No dropouts reported	• 12 men and 19 women • Mean age: 46 years	National Institutes of Health National Sunflower Association
Castro 2000 (81) Spain	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	9.4 8.6	24.3 24.8	4.3 4.7		Initial: 22, final: 21 Reason for loss: illness (n=1)	• 21 men • Mean age: 23 years	Investigaciones de la Seguridad Social Koype Co

u
3
_

Kris-Etherton 1999 (82) USA	Randomized crossover design with four interventions Experimental period: 24 days	1. 2. 3. 4.	6.7 6.7 6.7 7.7	11.5 20.2 16.3 17.3	5.8 5.8 8.6 9.6		Initial: 26, final: 22 Reason for loss: poor compliance (n=2), moved outside the area (n=2)	9 men and 13 women Mean age: 34 years	The Peanut Institute
Müller 2003 (83) Norway	Randomized crossover design with three interventions Experimental period: 22 days	1. 2. 3.	28.9 13.7 5.7	3.9 2.6 13.3	2.8 1.8 14.8	0.3 0.1 0.2	Initial: 31, final: 25 Reason for loss: not reported	• 25 women • Mean age: 31 years	The Norwegian Research Council Mills DA
Nielsen 2002 <i>(84)</i> Denmark	Randomized crossover design with three interventions Experimental period: 21 days	1. 2. 3.	10.5 11.5 11.5	14.5 16.9 7.6	6.5 2.3 11.7		Initial: 18, final: 18 No dropouts reported	• 18 men • Mean age: 24 years	Not reported
Poppitt 2002 (85) New Zealand	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	19.2 14.4	5.8 7.7	13.4 15.4		Initial: 20, final: 20 No dropouts reported	20 men Mean age: Not reported	New Zealand Dairy Board Auckland Uniservices Maurice & Phyllis Paykel Trust
Rajaram 2001 <i>(86)</i> USA	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	8.2 8.8	11.0 18.9	6.3 10.7		Initial: 24, final: 23 Reason for loss: not reported	14 men and 9 women Mean age: Not reported	National Pecan Sellers Association
Sanders 2003 (87) United Kingdom	Randomized crossover design with three interventions Experimental period: 14 days	1. ⁴ 2. 3.	9.1 9.8 10.5	11.8 19.9 12.3	5.9 6.3 6.1	9.2 0.1 0.1	Initial: 36, final: 29 Reason for loss: personal reasons (n=7)	29 men Mean age: 24 years	Ministry of Agriculture, Food and Fisheries The Medical Research Council
Wagner 2001 (88) Austria	Randomized crossover design with two interventions Experimental period: 14 days	1. 2.	8.5 8.4	9.8 14.5	11.5 6.9		Initial: 28, final: 28 No dropouts reported	28 men Mean age: 24 years	Not reported
Kratz 2002 (89) Germany	Randomized parallel design with three interventions Experimental period: 28 days	1. 2. 3.	9.1 10.7 10.0	19.3 23.3 8.7	9.0 3.4 18.5		Initial: 69, final: 58 Reason for loss: illness (n=6), poor compliance (n=5)	Diet 1: 10 men and 8 women Diet 2: 11 men and 9 women Diet 3: 10 men and 10 women Mean age: 26 years	Central Marketing Agency of the German Agricultural Industry The German Union for the Promotion of Oil and Protein Plants The Austrian Science Foundation The Brökelmann Ölmühle Company
Lichtenstein 2006 (90) USA	Randomized crossover design with five interventions Experimental period: 35 days	1. 2. 3. 4. 5. ⁴	6.5 4.9 5.8 6.8 7.3	6.3 6.1 18.8 6.7 8.1	12.3 14.1 2.3 13.2 8.1	0.6 0.6 0.3 0.5 2.4	Initial: 42 (including 10 replacers), final: 30 Reason for loss: time constraints (n=3), poor compliance (n=4), change in medical status (n=2), loss of medical insurance (n=1), moved out of the state (n=1), or dislike of the food (n=1)	 14 men and 16 women Mean age: 63 years 	The National Institutes of Health US Department of Agriculture
Motard-Belanger 2008 (91) Canada	Randomized crossover design with four interventions Experimental period: 28 days	1. 2. 3. ⁴ 4. ⁴	18.5 18.3 19.4 18.0	11.8 11.8 10.0 10.1	4.6 4.4 3.5 4.0	0.8 1.5 3.7 3.7	Initial: 48, final: 38 Reason for loss: not reported	38 men Mean age: 33 years	Dairy Farmers of Canada Novalait Inc Natural Sciences and Engineering Research Council of Canada
Rajaram 2009 <i>(92)</i> USA	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	9.4 8.0	9.4 8.0	4.3 10.8	1.0 0.8	Initial: 27, final: 25 Reason for loss: time constraints (n=2)	• 14 men and 11 women • Age: 23–65 years	California Walnut Commission

Reference and Study design		Composition ¹					Participants		
country	Study design	Diet	S	М	Р	T	Partic	ipants	Funding
Gillingham 2011 (93) Canada	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3.	11.2 5.6 6.1	16.1 22.9 15.9	6.5 5.7 12.3		Initial: 39, final: 36 Reason for loss: relocation of residence (n=2), work-related issues (n=1)	13 men and 23 women Mean age: 48 years	Flax Canada 2015 Canola Council of Canada Agri-Food Research & Development Initiative
Iggman 2011 <i>(94)</i> Sweden	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	19.6 7.9	11.1 17.4	3.9 9.6		Initial: 20, final: 20 No dropouts reported	14 men and 6 women Mean age: 51 years	Not reported
Marin 2011 (95) Spain	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	8.8 8.8	13.0 23.4	5.0 4.6		Initial: 59, final: 59 No dropouts reported	31 men and 28 women Mean age: 21 years	Ministerio de Ciencia e Innovacion / Spanish Ministry of Health CIBER Fisiopatologia de la Obesidad y Nutricion Consejeria de Innovacion Consejeria de Salud
Roussell 2012 (96) USA	Randomized crossover design with two interventions Experimental period: 35 days	1. 2.	6.0 6.0	9.0 11.0	8.0 7.0		Initial: 42, final: 36 Reason for loss: job change (n=1), illness (n=1), poor compliance (n=4)	15 men and 21 women Mean age: 50 years	Beef Checkoff Program National Institutes of Health
Zhao 2004 (97) USA	Randomized crossover design with three interventions Experimental period: 42 days	1. 2. 3.	12.7 8.5 8.2	13.2 12.2 12.3	8.7 16.4 17.2		Initial: 23, final: 23 No dropouts reported	• 20 men and 3 women • Mean age: 50 years	California Walnut Commission Walnut Marketing Board
Sabaté 2003 <i>(98)</i> USA	Randomized crossover design with three interventions Experimental period: 28 days	1. 2. 3.	8.2 8.0 7.7	12.1 16.5 19.4	6.2 7.5 8.7		Initial: 27, final: 25 Reason for loss: poor compliance (n=2)	• 14 men and 11 women • Mean age: 41 years	Almond Board of California
Curb 2000 (99) USA	Randomized crossover design with three interventions Experimental period: 30 days	1. 2. 3.	13.4 8.6 8.6	11.5 14.4 19.2	8.6 6.7 5.8		Initial: 34, final: 30 Reason for loss: not reported	• 15 men and 15 women • Age: 18-53 years	US Army Medical Research Acquisition Activity
Cater 2001 (100) USA	Randomized crossover design with two interventions Experimental period: 21 days	1. 2.	23.1 2.9	15.1 37.5	3.3 2.2		Initial: 7, final: 7 No dropouts reported	• 7 men • Mean age: 66 years	National Institutes of Health
Lacroix 2012 (101) Canada	Randomized crossover design with two interventions Experimental period: 28 days	1. 2.	9.9 10.3	14.2 12.8	5.9 5.8	0.6 1.8	Initial: 72, final: 61 Reason for loss: protocol too demanding (n=8), change of menopausal status (n=2), missing data (n=1)	61 women Mean age: 64 years	Dairy Farmers of Canada Dairy Australia Agriculture and Agri-Food Canada The Canadian Dairy Commission

S, saturated fatty acids; M, cis-monounsaturated fatty acids; P, cis-polyunsaturated fatty acids; T, trans-fatty acids

- ¹ The fatty acid composition of the diets is reported as a percentage of total energy intake
- ² Both publications reported data from the same study; the study is referred to as "Mensink 1987" in the table in risk of bias figures in **Annex 8**
- ³ Two separate studies were reported in a single publication; the studies are referred to as "Anderson 1976 A" and "Anderson 1976 B" in the table and risk of bias figures in **Annex 8**
- ⁴ Studies were not used, because the intake of *trans*-fatty acids was > 2% of energy
- ⁵ Both publications reported data from the same study; the study is referred to as "Lewis 1981" in the table and risk of bias figures in **Annex 8**
- ⁶ Both publications reported data from the same study; the study is referred to as "Wardlaw 1991" in the table and risk of bias figures in **Annex 8**
- Two separate studies were reported in a single publication; the studies are referred to as "Kris-Etherton 1993 A" and "Kris-Etherton 1993 B" in the table and risk of bias figures in Annex 8
- Both publications reported data from the same study; the study is referred to as "Judd 1988" in the table and risk of bias figures in Annex 8
- 9 All three publications reported data from the same study; the study is referred to as "Lichtenstein 1993" in the table and risk of bias figures in Annex 8
- Two separate studies were reported in a single publication; the studies are referred to as "Fielding 1995 A" and "Fielding 1995 B" in the table and risk of bias figures in **Annex 8**Both publications reported data from the same study; the study is referred to as "Judd 2002" in the table and risk of bias figures in **Annex 8**

Table 2. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from SFA is replaced isocalorically by carbohydrates, *cis*-MUFA or *cis*-PUFA

	Unit	Change	e per 1% of energy re	placed	
Lipid or lipoprotein		SFA → Carb	SFA → MUFA	SFA → PUFA	No¹
ΔTotal cholesterol	mmol/L	-0.041	-0.046	-0.064	177/74
95% Cl ²		-0.047 to -0.035	-0.051 to -0.040	-0.070 to -0.058	
P-value		<0.001	<0.001	<0.001	
ΔLDL cholesterol	mmol/L	-0.033	-0.042	-0.055	165/69
95% CI		-0.039 to -0.027	-0.047 to -0.037	-0.061 to -0.050	
P-value		<0.001	<0.001	<0.001	
ΔHDL cholesterol	mmol/L	-0.010	-0.002	-0.005	163/68
95% CI		-0.012 to -0.008	-0.004 to 0.000	-0.006 to -0.003	
P-value		<0.001	0.014	<0.001	
ΔTriglyceride	mmol/L	0.011	-0.004	-0.010	172/72
95% CI		0.007 to 0.014	-0.007 to -0.001	-0.014 to -0.007	
P-value		<0.001	0.022	<0.001	
ΔTotal to HDL cholesterol ratio		0.001	-0.027	-0.034	159/66
95% CI		-0.006 to 0.007	-0.033 to -0.022	-0.040 to -0.028	
P-value		0.842	<0.001	<0.001	
ΔLDL to HDL cholesterol ratio		-0.007	-0.029	-0.034	161/67
95% CI		-0.013 to -0.001	-0.034 to -0.024	-0.040 to -0.029	
P-value		0.017	<0.001	<0.001	
ΔTriglyceride to HDL cholesterol ratio		0.014	-0.002	-0.005	161/67
95% CI		0.010 to 0.018	-0.005 to 0.002	-0.009 to -0.002	
P-value		<0.001	0.342	0.004	
ΔΑροΑ-Ι	mg/dL	-7.0	-1.8	-4.9	102/41
95% CI		-9.0 to -5.1	-3.7 to 0.1	−7.3 to −2.5	
P-value		<0.001	0.064	<0.001	
ΔΑροΒ	mg/dL	-3.6	-7.8	-10.2	104/42
95% CI		−5.4 to −1.7	−9.5 to −6.0	-12.4 to -8.1	
P-value		<0.001	<0.001	<0.001	

 $ApoA-I, a polipoprotein A-I; Apo-B, a polipoprotein B; Carb, carbohydrates; CI, confidence interval; \emph{cis}-MUFA, \emph{cis}-monouns aturated fatty acids; \emph{cis}-PUFA, \emph{cis}-polyuns aturated fatty acids; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids$

¹ Number of diets/number of studies

² The 95% CIs refer to the regression coefficients on the line directly above

Table 3. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates, *cis*-MUFA or *cis*-PUFA is replaced isocalorically by SFA

	Unit	Change			
Lipid or lipoprotein		Carb → SFA	MUFA → SFA	PUFA → SFA	No ¹
ΔTotal cholesterol	mmol/L	0.045	0.049	0.066	177/74
95% Cl ²		0.038 to 0.051	0.043 to 0.055	0.060 to 0.073	
P-value		<0.001	<0.001	<0.001	
ΔLDL cholesterol	mmol/L	0.036	0.045	0.058	165/69
95% CI		0.030 to 0.043	0.039 to 0.051	0.052 to 0.064	
P-value		<0.001	<0.001	<0.001	
ΔHDL cholesterol	mmol/L	0.011	0.003	0.005	163/68
95% CI		0.010 to 0.013	0.001 to 0.004	0.004 to 0.007	
P-value		<0.001	0.001	<0.001	
ΔTriglyceride	mmol/L	-0.012	0.004	0.010	172/72
95% CI		-0.015 to -0.008	0.000 to 0.007	0.006 to 0.014	
P-value		<0.001	0.041	<0.001	
ΔTotal to HDL cholesterol ratio		-0.002	0.028	0.034	159/66
95% CI		-0.009 to 0.005	0.021 to 0.034	0.027 to 0.041	
P-value		0.553	<0.001	<0.001	
ΔLDL to HDL cholesterol ratio		0.007	0.030	0.035	161/67
95% CI		0.001 to 0.014	0.024 to 0.036	0.028 to 0.041	
P-value		0.033	<0.001	<0.001	
ΔTriglyceride to HDL cholesterol ratio		-0.016	0.001	0.004	161/67
95% CI		-0.020 to -0.012	-0.003 to 0.004	0.001 to 0.008	
P-value		<0.001	0.680	0.026	
ΔΑροΑ-Ι	mg/dL	8.4	2.7	6.3	102/41
95% CI		6.4 to 10.5	0.7 to 4.8	3.9 to 8.7	
P-value		<0.001	0.008	<0.001	
ΔΑροΒ	mg/dL	3.7	8.1	10.3	104/42
95% CI		1.7 to 5.8	6.1 to 10.1	7.7 to 12.8	
P-value		0.001	<0.001	<0.001	

 $ApoA-I, a polipoprotein A-I; Apo-B, a polipoprotein B; Carb, carbohydrates; CI, confidence interval; \emph{cis}-MUFA, \emph{cis}-monouns aturated fatty acids; \emph{cis}-PUFA, \emph{cis}-polyuns aturated fatty acids; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids$

¹ Number of diets/number of studies

² The 95% CIs refer to the regression coefficients on the line directly above

Table 4. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by SFA: impact of baseline levels

Lipid or lipoprotein	Unit	Change per 1% of energy replaced	No	Change per 1% of energy replaced	No¹	
		Below median ²		Above median		
ΔTotal cholesterol	mmol/L	0.035	82/37	0.050	95/37	
95% CI ³		0.023 to 0.048		0.043 to 0.057		
P-value		<0.001		<0.001		
ΔLDL cholesterol	mmol/L	0.029	79/35	0.041	86/34	
95% CI		0.020 to 0.039		0.032 to 0.049		
P-value		<0.001		<0.001		
ΔHDL cholesterol	mmol/L	0.008	81/34	0.013	82/34	
95% CI		0.005 to 0.011		0.011 to 0.016		
P-value		<0.001		<0.001		
ΔTriglyceride	mmol/L	-0.011	83/36	-0.013	89/36	
95% CI		−0.015 to −0.006		-0.019 to -0.007		
P-value		<0.001		<0.001		
ΔTotal to HDL cholesterol ratio		0.002	76/33	-0.006	83/33	
95% CI		-0.008 to 0.012		-0.016 to 0.004		
P-value		0.695		0.246		
ΔLDL to HDL cholesterol ratio		0.007	78/34	0.006	83/33	
95% CI		-0.002 to 0.016		-0.004 to 0.017		
P-value		0.103		0.218		
ΔTriglyceride to HDL cholesterol ratio		-0.012	78/34	-0.019	83/33	
95% CI		-0.016 to -0.008		-0.026 to -0.013		
P-value		<0.001		<0.001		

CI, confidence interval; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids

¹ Number of diets/number of studies

Number of diets/number of studies
 The median level when subjects consumed a standardized fat-free diet was for total cholesterol 4.45 mmol/L, for LDL cholesterol 2.89 mmol/L, for HDL cholesterol 0.97 mmol/L, for triglyceride 1.48 mmol/L, for the total to HDL cholesterol ratio 4.36, for the LDL cholesterol to HDL cholesterol ratio 2.76 and for the for the triglyceride to HDL cholesterol ratio 1.36
 The 95% CIs refer to the regression coefficients on the line directly above

Table 5. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by SFA: impact of gender

A total contraction	11.24	Change per 1% of energy replaced	N.	Change per 1% of energy replaced	No¹	
Lipid or lipoprotein	Unit	Men only	No	Women only or men and women		
ΔTotal cholesterol	mmol/L	0.037	85/38	0.049	92/36	
95% Cl ²		0.028 to 0.047		0.041 to 0.057		
P-value		<0.001		<0.001		
ΔLDL cholesterol	mmol/L	0.026	73/33	0.041	92/36	
95% CI		0.014 to 0.038		0.033 to 0.048		
P-value		<0.001		<0.001		
ΔHDL cholesterol	mmol/L	0.007	71/32	0.013	92/36	
95% CI		0.004 to 0.010		0.011 to 0.016		
P-value		<0.001		<0.001		
ΔTriglyceride	mmol/L	-0.014	82/37	-0.010	90/35	
95% CI		-0.020 to -0.008		-0.015 to -0.005		
P-value		<0.001		<0.001		
ΔTotal to HDL cholesterol ratio		0.002	71/32	-0.004	88/34	
95% CI		-0.014 to 0.018		-0.011 to 0.003		
P-value		0.808		0.216		
ΔLDL to HDL cholesterol ratio		0.008	71/32	0.007	90/35	
95% CI		-0.006 to 0.021		-0.001 to 0.014		
P-value		0.269		0.088		
ΔTriglyceride to HDL cholesterol ratio		-0.012	71/32	-0.017	90/35	
95% CI		-0.020 to -0.005		-0.022 to -0.012		
P-value		0.002		<0.001		

 ${\sf CI, confidence\,interval; HDL, high-density\,lipoprotein; LDL, low-density\,lipoprotein; SFA, saturated\,fatty\,acids}$

¹ Number of diets/number of studies

² The 95% CIs refer to the regression coefficients on the line directly above

Table 6. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by SFA: impact of year of publication

Lipid or lipoprotein	Unit	Change per 1% of energy replaced	No	Change per 1% of energy replaced	No¹	
Lipid or lipoprotein	Unit	Published before 1993	NO	Published in 1993 or later		
ΔTotal cholesterol	mmol/L	0.045	77/34	0.045	100/40	
95% CI ²		0.035 to 0.054		0.036 to 0.054		
P-value		<0.001		<0.001		
ΔLDL cholesterol	mmol/L	0.035	69/31	0.038	96/38	
95% CI		0.024 to 0.046		0.030 to 0.046		
P-value		<0.001		<0.001		
ΔHDL cholesterol	mmol/L	0.011	67/30	0.012	96/38	
95% CI		0.007 to 0.014		0.010 to 0.014		
P-value		<0.001		<0.001		
ΔTriglyceride	mmol/L	-0.014	72/32	-0.010	100/40	
95% CI		-0.019 to -0.009		-0.015 to -0.004		
P-value		<0.001		0.002		
ΔTotal to HDL cholesterol ratio		0.003	65/29	-0.004	94/37	
95% CI		-0.008 to 0.015		-0.013 to 0.005		
P-value		0.543		0.344		
ΔLDL to HDL cholesterol ratio		0.010	65/29	0.007	96/38	
95% CI		0.000 to 0.020		-0.002 to 0.015		
P-value		0.048		0.143		
ΔTriglyceride to HDL cholesterol ratio		-0.014	65/29	-0.017	96/38	
95% CI		-0.020 to -0.009		-0.023 to -0.012		
P-value		<0.001		<0.001		

 ${\it CI, confidence interval; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids}$

 $^{^1}$ $\,$ Number of diets/number of studies 2 $\,$ The 95% CIs refer to the regression coefficients on the line directly above

Table 7. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by SFA: exclusion of studies using liquid formula diets

		Change per 1% of energy replaced		
Lipid or lipoprotein	Unit	Carb → SFA	No¹	
ΔTotal cholesterol	mmol/L	0.046	166/69	
95% Cl ²		0.039 to 0.052		
P-value		<0.001		
ΔLDL cholesterol	mmol/L	0.037	154/64	
95% CI		0.031 to 0.044		
P-value		<0.001		
ΔHDL cholesterol	mmol/L	0.011	152/63	
95% CI		0.010 to 0.013		
P-value		<0.001		
ΔTriglyceride	mmol/L	-0.012	163/68	
95% CI		-0.016 to -0.008		
P-value		0.001		
ΔTotal to HDL cholesterol ratio		-0.002	150/62	
95% CI		-0.009 to 0.004		
P-value		0.485		
ΔLDL to HDL cholesterol ratio		0.007	152/63	
95% CI		0.000 to 0.014		
P-value		0.040		
ΔTriglyceride to HDL cholesterol ratio		-0.016	152/63	
95% CI		-0.020 to -0.012		
P-value		<0.001		
ΔΑροΑ-Ι	mg/dL	8.4	100/40	
95% CI		6.4 to 10.5		
P-value		<0.001		
ΔΑροΒ	mg/dL	3.7	102/41	
95% CI		1.6 to 5.8		
P-value		0.001		

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; Carb, carbohydrates; CI, confidence interval; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids

¹ Number of diets/number of studies

 $^{^{2}\,\,}$ The 95% CIs refer to the regression coefficients on the line directly above

Table 8. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by lauric acid (C12:0), myristic acid (C14:0), palmitic acid (C16:0) or stearic acid (C18:0)

			Change per 1% of	energy replaced		
Lipid or lipoprotein	Unit	Carb → C12:0	Carb → C14:0	Carb → C16:0	Carb → C18:0	No ¹
ΔTotal cholesterol	mmol/L	0.029	0.060	0.041	-0.006	134/52
95% Cl ²		0.014 to 0.045	0.042 to 0.077	0.030 to 0.052	-0.019 to 0.007	
P-value		<0.001	<0.001	<0.001	0.384	
ΔLDL cholesterol	mmol/L	0.017	0.044	0.036	-0.003	130/50
95% CI		0.003 to 0.031	0.028 to 0.060	0.026 to 0.046	-0.015 to 0.009	
P-value		0.019	<0.001	<0.001	0.606	
ΔHDL cholesterol	mmol/L	0.019	0.021	0.010	0.000	132/51
95% CI		0.016 to 0.023	0.017 to 0.025	0.007 to 0.013	-0.003 to 0.003	
P-value		<0.001	<0.001	<0.001	0.853	
ΔTriglyceride	mmol/L	-0.015	-0.011	-0.011	-0.005	135/53
95% CI		-0.023 to -0.007	-0.020 to -0.002	-0.017 to -0.006	-0.012 to 0.001	
P-value		<0.001	0.018	<0.001	0.110	
ΔTotal to HDL cholesterol ratio		-0.035	-0.009	0.006	-0.002	125/48
95% CI		-0.048 to -0.022	-0.023 to 0.006	-0.003 to 0.015	-0.013 to 0.009	
P-value		<0.001	0.244	0.180	0.676	
ΔLDL to HDL cholesterol ratio		-0.024	0.000	0.013	-0.001	130/50
95% CI		-0.036 to -0.013	-0.013 to 0.014	0.005 to 0.021	-0.011 to 0.009	
P-value		<0.001	0.941	0.002	0.831	
ΔTriglyceride to HDL cholesterol ratio		-0.024	-0.018	-0.015	-0.003	131/51
95% CI		-0.032 to -0.017	-0.027 to -0.010	-0.020 to -0.009	-0.009 to 0.004	
P-value		<0.001	<0.001	<0.001	0.407	
ΔΑροΑ-Ι	mg/dL	19.2	6.8	6.5	-1.4	88/34
95% CI		14.6 to 23.7	0.5 to 13.1	3.8 to 9.3	-4.4 to 1.7	
P-value		<0.001	0.034	<0.001	0.374	
ΔΑροΒ	mg/dL	-1.3	2.0	3.3	-1.8	91/35
95% CI		-6.8 to 4.2	-2.9 to 6.9	-0.2 to 6.9	-5.7 to 2.1	
P-value		0.627	0.417	0.065	0.368	

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; Carb, carbohydrates; CI, confidence interval; cis-MUFA, cis-monounsaturated fatty acids; cis-PUFA, cis-polyunsaturated fatty acids; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids

 $^{^{\}scriptscriptstyle 1} \ \ {\hbox{Number of diets/number of studies}}$

 $^{^{2}\,\,}$ The 95% CIs refer to the regression coefficients on the line directly above

Table 9. Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by lauric acid (C12:0), myristic acid (C14:0), palmitic acid (C16:0) or stearic acid (C18:0): exclusion of studies using liquid formula diets

			Change per 1% of	energy replaced		N-1
Lipid or lipoprotein	Unit	Carb → C12:0	Carb → C14:0	Carb → C16:0	Carb → C18:0	No¹
ΔTotal cholesterol	mmol/L	0.033	0.058	0.041	-0.003	122/48
95% CI ²		0.014 to 0.052	0.039 to 0.076	0.028 to 0.053	-0.017 to 0.012	
P-value		0.001	<0.001	<0.001	0.717	
ΔLDL cholesterol	mmol/L	0.019	0.042	0.036	0.000	118/46
95% CI		0.002 to 0.036	0.026 to 0.059	0.025 to 0.048	-0.013 to 0.013	
P-value		0.031	<0.001	<0.001	0.977	
ΔHDL cholesterol	mmol/L	0.021	0.020	0.010	-0.001	120/47
95% CI		0.017 to 0.026	0.016 to 0.025	0.007 to 0.012	-0.004 to 0.003	
P-value		<0.001	<0.001	<0.001	0.684	
ΔTriglyceride	mmol/L	-0.016	-0.011	-0.012	-0.005	123/49
95% CI		-0.025 to -0.006	-0.020 to -0.002	-0.018 to -0.006	-0.012 to 0.002	
P-value		0.001	0.023	<0.001	0.177	
ΔTotal to HDL cholesterol ratio		-0.033	-0.009	0.006	0.002	116/45
95% CI		-0.048 to -0.019	-0.023 to 0.006	-0.003 to 0.016	-0.010 to 0.013	
P-value		<0.001	0.223	0.188	0.779	
ΔLDL to HDL cholesterol ratio		-0.022	0.000	0.013	0.003	118/46
95% CI		-0.035 to -0.009	-0.013 to 0.013	0.005 to 0.022	-0.007 to 0.013	
P-value		0.001	0.994	0.003	0.604	
ΔTriglyceride to HDL cholesterol ratio		-0.025	-0.018	-0.014	-0.002	120/47
95% CI		-0.034 to -0.016	-0.026 to -0.009	-0.020 to -0.008	-0.009 to 0.005	
P-value	P-value		<0.001	<0.001	0.539	


 ${\sf CI, confidence\,interval; HDL, high-density\,lipoprotein; LDL, low-density\,lipoprotein; SFA, saturated\,fatty\,acids}$

¹ Number of diets/number of studies

² The 95% CIs refer to the regression coefficients on the line directly above

7. Figures

Figure 1. Flow diagram of study selection



ANNEX 1.

PICO questions

- 1. What is the effect in the population of reduced percentage of total energy intake from saturated fatty acids (SFA) relative to higher intake for reduction in risk of noncommunicable diseases (NCDs)?
- 2. What is the effect in the population of consuming less than 10% of total energy as SFA relative to more than 10% total energy as SFA for reduction in risk of NCDs?
- 3. What is the effect in the population of a reduction in percentage of total energy intake from SFA from 10% in gradual increments relative to higher intake for reduction in risk of NCDs?
- 4. What is the effect in the population of reduced percentage of total energy intake from long-chain SFA, very long-chain SFA and medium-chain SFA relative to higher intake for reduction in risk of NCDs?
- 5. What is the effect in the population of reduced percentage of total energy intake from lauric acid, myristic acid, palmitic acid or stearic acid relative to higher intake for reduction in risk of NCDs?
- 6. What is the effect in the population of replacing SFA with carbohydrates (refined vs. unrefined), *cis*-monounsaturated fatty acids (*cis*-MUFA), *cis*-polyunsaturated fatty acids (*cis*-PUFA), protein or trans-fatty acids (TFA) relative to no replacement for reduction in risk of NCDs?

ANNEX 2.

Priority outcomes

- 1. All-cause mortality
- 2. Coronary heart disease (CHD) incidence, CHD mortality, and CHD morbidity
- 3. Cardiovascular disease (CVD) incidence (as a composite indicator defined by study authors), CVD mortality, and CVD morbidity
- 4. Stroke including stroke incidence (type of stroke), stroke mortality, and stroke morbidity
- 5. Blood lipids including total cholesterol, low-density lipoprotein (LDL) cholesterol, high-density lipoprotein (HDL) cholesterol, triglyceride, LDL cholesterol to HDL cholesterol ratio, total cholesterol to HDL cholesterol ratio, triglyceride to HDL cholesterol ratio and lipoprotein (a)
- 6. Adverse effects reported by study authors

ANNEX 3.

Search strategy

PubMed

((((((("comparative study"[Publication Type]) OR "randomized controlled trial"[Publication Type]) OR "controlled clinical trial"[Publication Type]))

AND

(((("cholesterol/blood"[MeSHTerms])OR"cholesterol, Idl/blood"[MeSHTerms])OR"lipids/

AND

"humans"[MeSH Terms]))


AND

((dietary fat*[MeSH Terms]) OR (((((palmitic acid*[MeSH Terms]) OR stearic acid*[MeSH Terms])) OR myristic acid*[MeSH Terms])) OR lauric acid*[MeSH Terms]))

ANNEX 4.

Residuals analysis

Scatterplot of the relationship between SFA intake (model 1 in **Section 2.2.4**) and the difference between observed and predicted serum LDL cholesterol concentrations (residuals). Each point refers to one of the 165 diets from the 69 studies as used for the calculations (see **Table 1**). "Predicted" values were calculated as the intrinsic level of the group under study plus the predicted change induced by the experimental diet.




SFA, saturated fatty acids; LDL, low-density lipoprotein

ANNEX 5.

Relationship between observed and predicted serum LDL cholesterol concentrations

Each point refers to one of the 165 diets from the 69 studies as used for the calculations in regression model 1 (see **Section 2.2.4 and Table 2**). "Predicted" values were calculated as the intrinsic level of the group under study plus the predicted change induced by their experimental diet



ANNEX 6.

Results of cis-PUFA replacement

Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from *cis*-PUFA in the diet is replaced isocalorically by carbohydrates, SFA or *cis*-MUFA.

		Chan	ge per 1% of energy rep	laced	No.1
Lipid or lipoprotein	Unit	PUFA → Carb	PUFA → SFA	PUFA → MUFA	No ¹
ΔTotal cholesterol	mmol/L	0.019	0.066	0.016	177/74
95% CI ²		0.013 to 0.025	0.060 to 0.073	0.011 to 0.022	
P-value		<0.001	<0.001	<0.001	
ΔLDL cholesterol	mmol/L	0.019	0.058	0.012	165/69
95% CI		0.012 to 0.025	0.052 to 0.064	0.007 to 0.017	
P-value		<0.001	<0.001	<0.001	
ΔHDL cholesterol	mmol/L	-0.005	0.005	0.003	163/68
95% CI		-0.007 to -0.004	0.004 to 0.007	0.001 to 0.004	
P-value		<0.001	<0.001	<0.001	
ΔTriglyceride	mmol/L	0.020	0.010	0.006	172/72
95% CI		0.016 to 0.024	0.006 to 0.014	0.003 to 0.009	
P-value		<0.001	<0.001	<0.001	
ΔTotal to HDL cholesterol ratio		0.032	0.034	0.005	159/66
95% CI		0.025 to 0.039	0.027 to 0.041	0.000 to 0.011	
P-value		<0.001	<0.001	0.053	
ΔLDL to HDL cholesterol ratio		0.024	0.035	0.004	161/67
95% CI		0.017 to 0.031	0.028 to 0.041	-0.001 to 0.009	
P-value		<0.001	<0.001	0.104	
ΔTriglyceride to HDL cholesterol ratio		0.018	0.004	0.003	161/67
95% CI		0.014 to 0.022	0.001 to 0.008	0.000 to 0.006	
P-value		<0.001	0.026	0.040	
ΔΑροΑ-Ι	mg/dL	-1.8	6.3	3.4	104/42
95% CI		-4.0 to 0.3	3.9 to 8.7	1.6 to 5.3	
P-value		0.097	<0.001	0.001	
ΔΑροΒ	mg/dL	5.7	10.3	1.8	102/41
95% CI		3.3 to 8.1	7.7 to 12.8	-0.2 to 3.8	
P-value		<0.001	<0.001	0.074	

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; Carb, carbohydrates; CI, confidence interval; cis-MUFA, cis-monounsaturated fatty acids; cis-PUFA, cis-polyunsaturated fatty acids; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids

¹ Number of diets/number of studies

The 95% CIs refer to the regression coefficients on the line directly above

ANNEX 7.

Results of cis-MUFA replacement

Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from *cis*-MUFA is replaced isocalorically by carbohydrates, SFA or by *cis*-PUFA.

		Chan	ge per 1% of energy rep	laced	
Lipid or lipoprotein	Unit	MUFA → Carb	MUFA → SFA	MUFA → PUFA	No¹
ΔTotal cholesterol	mmol/L	0.003	0.049	-0.018	177/74
95% Cl ²		-0.002 to 0.008	0.043 to 0.055	-0.023 to -0.013	
P-value		0.227	<0.001	<0.001	
ΔLDL cholesterol	mmol/L	0.007	0.045	-0.013	165/69
95% CI		0.002 to 0.012	0.039 to 0.051	-0.018 to -0.009	
P-value		0.012	<0.001	<0.001	
ΔHDL cholesterol	mmol/L	-0.008	0.003	-0.002	163/68
95% CI		-0.009 to -0.006	0.001 to 0.004	-0.004 to -0.001	
P-value		<0.001	0.001	0.002	
ΔTriglyceride	mmol/L	0.014	0.004	-0.007	172/72
95% CI		0.011 to 0.018	0.000 to 0.007	-0.010 to -0.004	
P-value		<0.001	0.041	<0.001	
ΔTotal to HDL cholesterol		0.026	0.028	-0.008	159/66
95% CI		0.020 to 0.032	0.021 to 0.034	-0.013 to -0.002	
P-value		<0.001	<0.001	0.005	
Δ LDL cholesterol to HDL cholesterol		0.020	0.030	-0.006	161/67
95% CI		0.014 to 0.026	0.024 to 0.036	-0.011 to -0.001	
P-value		<0.001	<0.001	0.018	
ΔTriglyceride to HDL cholesterol		0.015	0.001	-0.004	161/67
95% CI		0.012 to 0.019	-0.003 to 0.004	-0.007 to -0.001	
P-value		<0.001	0.680	0.009	
ΔΑροΑ-Ι	mg/dL	-5.0	2.7	-3.0	102/41
95% CI		-6.9 to -3.1	0.7 to 4.8	-5.0 to-1.0	
P-value		<0.001	0.008	0.004	
ΔΑροΒ	mg/dL	3.7	8.1	-2.7	104/42
95% CI		1.7 to 5.7	6.1 to 10.1	-4.7to -0.8	
P-value		0.001	<0.001	0.007	

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; Carb, carbohydrates; CI, confidence interval; cis-MUFA, cis-monounsaturated fatty acids; cis-PUFA, cis-polyunsaturated fatty acids; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids

¹ Number of diets/number of studies

² The 95% CIs refer to the regression coefficients on the line directly above

ANNEX 8.

Results of carbohydrate replacement

Estimated multiple regression equations for the mean changes in serum lipids and lipoproteins when 1% of energy in the diet from carbohydrates is replaced isocalorically by SFA, *cis*-MUFA or *cis*-PUFA.

		Chan	ge per 1% of energy rep	laced	
Lipid or lipoprotein	Unit	Carb → SFA	Carb → MUFA	Carb → PUFA	No ¹
ΔTotal cholesterol	mmol/L	0.045	-0.004	-0.022	177/74
95% CI ²		0.038 to 0.051	-0.010 to 0.001	-0.028 to -0.016	
P-value		<0.001	0.097	<0.001	
ΔLDL cholesterol	mmol/L	0.036	-0.009	-0.022	165/69
95% CI		0.030 to 0.043	-0.014 to -0.003	-0.028 to -0.015	
P-value		<0.001	0.002	<0.001	
ΔHDL cholesterol	mmol/L	0.011	0.008	0.006	163/68
95% CI		0.010 to 0.013	0.007 to 0.010	0.004 to 0.008	
P-value		<0.001	<0.001	<0.001	
ΔTriglyceride	mmol/L	-0.012	-0.015	-0.021	172/72
95% CI		-0.015 to -0.008	-0.018 to -0.011	-0.025 to -0.017	
P-value		<0.001	<0.001	<0.001	
ΔTotal to HDL cholesterol		-0.002	-0.029	-0.036	159/66
95% CI		-0.009 to 0.005	-0.035 to -0.023	-0.043 to -0.029	
P-value		0.553	<0.001	<0.001	
ΔLDL cholesterol to HDL cholesterol		0.007	-0.022	-0.027	161/67
95% CI		0.001 to 0.014	-0.028 to -0.016	-0.034 to -0.021	
P-value		0.033	<0.001	<0.001	
ΔTriglyceride to HDL cholesterol		-0.016	-0.016	-0.020	161/67
95% CI		-0.020 to -0.012	-0.020 to -0.013	-0.024 to -0.016	
P-value		<0.001	<0.001	<0.001	
ΔΑροΑ-Ι	mg/dL	8.4	5.5	2.3	104/42
95% CI		6.4 to 10.5	3.7 to 7.3	0.1 to 4.6	
P-value		<0.001	<0.001	0.042	
ΔΑροΒ	mg/dL	3.7	-4.4	-6.9	102/41
95% CI		1.7 to 5.8	-6.3 to -2.4	-9.1 to -4.6	
P-value		0.001	<0.001	<0.001	

 $ApoA-I, a polipoprotein A-I; Apo-B, a polipoprotein B; Carb, carbohydrates; CI, confidence interval; \emph{cis}-MUFA, \emph{cis}-monouns aturated fatty acids; \emph{cis}-PUFA, \emph{cis}-polyuns aturated fatty acids; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SFA, saturated fatty acids$

¹ Number of diets/number of studies

 $^{^{\}rm 2}$ $\,$ The 95% CIs refer to the regression coefficients on the line directly above

ANNEX 9.

Risk of bias assessment

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants (performance bias)	Blinding of personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Anderson 1976 A	•	?	•	?	•	•	•	•
Anderson 1976 B	•	?	•	?	•	•	•	•
Barr 1992	•	?	•	?	•	•	•	•
Becker 1983	•	?	•	?	•	•	•	•
Berglund 2007	•	?	•	?	•	•	•	•
Berry 1991	•	?	•	?	•	•	•	•
Berry 1992	•	?	•	?	•	•	•	•
Binkoski 2005	•	?	•	•	•	•	•	•
Bonanome 1988	•	?	•	?	•	•	•	•
Bonanome 1992	•	?	•	?	•	•	•	•
Brussaard 1980	?	?	•	?	•	•	•	•
Brussaard 1982	?	?	•	?	•	•	•	•
Castro 2000	•	?	•	•	•	•	•	•
Cater 1997	•	?	•	?	•	•	•	•
Cater 2001	•	?	•	?	•	•	•	•
Chan 1991	•	?	•	?	•	•	•	•
Curb 2000	•	?	•	?	•	•	•	•
Denke 1992	•	?	•	?	•	•	•	•
Dougherty 1995	•	?	•	?	•	•	•	•
Fielding 1995 A	•	?	•	?	•	•	•	•
Fielding 1995 B	•	?	•	?	•	•	•	•
Gillingham 2011	•	?	•	?	•	•	•	•
Ginsberg 1990	•	?	•	•	•	•	•	•
Ginsberg 1994	•	?	•	?	•	•	•	•
Ginsberg 1998	•	?	•	?	•	•	•	•
Grande 1972	•	?	•	?	•	•	•	•
Grundy 1986 A	•	?	•	?	•	•	•	•
Grundy 1986 B	•	?	•	?	•	•	•	•
Grundy 1988	•	?	•	?	•	•	•	•
Harris 1983	•	?	•	?	•	•	•	•
Howard 1995	•	?	•	•	•	•	•	•
Hunter 2000	•	?	•	?	•	•	•	•
lacono 1991	•	?	•	?	?	•	•	•
Iggman 2011	•	?	•	?	•	•	•	•
Judd 1988	•	?	•	?	•	•	•	•
Judd 1994	•	?	•	•	•	•	•	•
Judd 2002	•	?	•	?	•	•	•	•
Katan 1988	•	?	•	?	•	•	•	•
Kratz 2002	•	?	•	?	•	•	•	•
Kris-Etherton 1993 A	•	?	•	?	•	•	•	•
Kris-Etherton 1993 B	•	?	•	?	•	•	•	•
Kris-Etherton 1999	•	?	•	?	•	•	•	•

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants (performance bias)	Blinding of personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Lacroix 2012	•	?	•	?	•	•	•	•
Laine 1982	•	?	•	?	•	•	•	•
Lewis 1981	•	?	•	?	•	•	•	•
Lichtenstein 1993	•	?	•	?	•	•	•	•
Lichtenstein 1999	•	?	•	•	•	•	•	•
Lichtenstein 2006	•	?	•	•	•	•	•	•
Lovejoy 2002	•	?	•	?	•	•	•	•
Marckmann 1992	•	?	•	?	•	•	•	•
Marin 2011	•	?	9	?	0	•	•	•
Mattson 1985	•	?	•	?	•	•	•	•
Mazier 1997 McDonald 1989	•	?	•	?	•	•	•	•
Mensink 1987	•	?	•	?	•	•	•	•
Mensink 1989	?	•	?	•	?	•	•	•
Mensink 1990	•	?	•	?	•	•	•	•
Motard-Belanger 2008	?	•	?	•	•	•	•	•
Müller 1998	•	?	•	?	•	•	•	•
Müller 2003	•	?	•	?	•	•	•	•
Nielsen 2002	•	?	•	?	•	•	•	•
Park 1996	•	?	•	?	•	•	•	•
Poppitt 2002	?	•	?	•	•	•	•	•
Rajaram 2001	•	?	•	•	•	•	•	•
Rajaram 2009	•	?	•	?	•	•	•	•
Reiser 1985	•	?	•	?	•	•	•	•
Roussell 2012	•	?	•	?	•	•	•	•
Sabaté 2003	•	?	•	?	•	•	•	•
Sanders 2003	•	?	•	?	•	•	•	•
Sundram 1994	•	?	•	?	•	•	•	•
Sundram 1995	•	?	•	?	•	•	•	•
Thoistrup 1994	•	?	•	?	•	•	•	•
Tholstrup 1994 B	•	?	•	?	•	•	•	•
Thoistrup 1998	0	?	•	?	0	•	•	•
Valsta 1992	•	?	•	•	•	•	•	•
Vega-López 2006	•	?	•	?	•	•	•	•
Wagner 2001 Wahrburg 1992	•	?	•	?	?	•	•	•
Wardlaw 1990	•	?	•	•	•	•	•	•
Wardlaw 1990 Wardlaw 1991	•	?	•	•	•	•	•	•
Wolf 1983	•	?	•	?	•	•	•	•
Zhao 2004	•	?	•	?	•	•	•	•
Zock 1992	•	?	•	?	•	•	•	•
Zock 1994	•	?	•	?	•	•	•	•
	_	_						

- low risk of biasunclear risk of bias
- high risk of bias

GRADE evidence profiles

GRADE evidence profile 1

Question: What is the effect of replacing saturated fatty acids in the diet of adults with *cis*-polyunsaturated fatty acids?¹

Population: General adult population

			Quality assessment				No. of	Effect ⁹		
No. of studies ²	Design	Risk of bias ³	Inconsistency⁴	Indirectness⁵	Imprecision ⁶	Other considerations ⁷	participants ⁸	(95% CI)	Quality	Importance
Total cholesterol	(follow-up 13–91 d	ays; units mmol/L բ	oer 1% energy exch	ange; better indica	ted by lower values)					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.064 (-0.070, -0.058)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	er 1% energy exchai	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.055 (-0.061, -0.050)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (follow-up 13–91 da	ys; units mmol/L p	er 1% energy excha	nge; better indicate	ed by higher values)					
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.005 (-0.006, -0.003)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; เ	units mmol/L per 19	% energy exchange;	better indicated by	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.010 (-0.014, -0.007)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (follow-up 13	–91 days; unitless; l	petter indicated by	lower values)					
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	-0.034 (-0.040, -0.028)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol r	atio (follow-up 13–	91 days; unitless; be	etter indicated by lo	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.034 (-0.040, -0.029)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	rvalues)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.005 (-0.009, -0.002)	⊕⊕⊕⊕ HIGH	IMPORTANT
ApoA-I (follow-up	14–91 days; units	mg/dL per 1% ener	gy exchange; better	indicated by highe	er values)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	-4.9 (-7.3, -2.5)	⊕⊕⊕⊕ HIGH	IMPORTANT
Apo-B (follow-up	14–91 days; units r	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-10.2 (-12.4, -8.1)	⊕⊕⊕⊕ HIGH	IMPORTANT

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; CI, confidence interval; LDL, low-density lipoprotein; HDL, high-density lipoprotein; RCT, randomized controlled trial

- ¹ A mixture of dietary saturated fatty acids (SFA) was replaced with cis-polyunsaturated fatty acids (cis-PUFA) consisting of linoleic acid plus α-linolenic acid.
- ² Number of data points (diet groups) are provided in parentheses. Each data point contains mean dietary information on SFA, *cis*-PUFA and *cis*-monounsaturated fatty acid (*cis*-MUFA) intake as well as an associated change in a given serum lipid or lipoprotein for each study group (i.e. intervention and control groups) at the end of a dietary treatment period, and was extracted for all treatment groups within studies included in the analysis.
- 3 All studies were strictly controlled, dietary trials lasting from 13 to 91 days in which protein and cholesterol intakes were held constant. Some of the studies with parallel design were assessed as having unclear risk of bias in terms of randomization as the randomization procedure was not described adequately. Studies with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have an significant, systematic effect on study results. Blinding was not deemed to be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some studies may have been able to distinguish between intervention and control diets, this was not expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were judged as having a low risk of bias.
- This analysis was conducted as a multiple regression in which data points (see Footnote 2) were directly extracted from each study, rather than extraction of mean differences between groups within each included study. As a result, directly measuring between-study variability in a quantitative manner is not feasible. Qualitative assessment of the included studies and the strength and consistency of the results of the multiple regression, however, indicate that any inconsistency is unlikely to decrease confidence in the results of the multiple regression analysis and is therefore not considered to be serious.
- 5 All studies directly assessed the effect of modifying dietary fat intake such that models could be derived, which provide estimates of the effects of exchanging SFA with other nutrients on serum lipids and lipoproteins, which were priority health outcomes decided upon prior to initiating review. All studies were conducted in the population of interest (adults without disturbances of lipid metabolism or diabetes).
- ⁶ Imprecision was assessed using the 95% CI of the regression coefficient as a proxy for the 95% CI of a pooled estimate of effect, the rationale being that the regression coefficient is a direct measure of the effect of reducing SFA intake on a particular serum lipid or lipoprotein and the 95% CI is a measure of variability of that effect.. Unless otherwise noted, the 95% CI does not cross a threshold of irrelevant benefit or important harm and the outcome has not been downgraded for serious imprecision.
- ⁷ Publication bias was not formally assessed.
- ⁸ Total number of participants. Data points containing dietary information on SFA, *cis*-PUFA and *cis*-MUFA intake as well as an associated change in a given serum lipid or lipoprotein were directly extracted from all study groups from included trials without distinction between intervention and control groups.
- The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of cis-PUFA.

GRADE evidence profile 2

Question: What is the effect of replacing saturated fatty acids in the diet of adults with cis-monounsaturated fatty acids?¹

Population: General adult population

			Quality assessment				No. of	F# - +49		
No. of studies ²	Design	Risk of bias ³	Inconsistency⁴	Indirectness ⁵	Imprecision ⁶	Other considerations ⁷	No. of participants ⁸	Effect ⁹ (95% CI)	Quality	Importance
Total cholesterol	(follow-up 13–91 d	ays; units mmol/L p	oer 1% energy exch	ange; better indica	ted by lower values					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.046 (-0.051, -0.040)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	er 1% energy exchai	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.042 (-0.047, -0.037)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (follow-up 13–91 da	ays; units mmol/L p	er 1% energy excha	nge; better indicate	ed by higher values)					
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.002 (-0.004, 0.000)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; เ	units mmol/L per 19	% energy exchange;	better indicated b	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.004 (-0.007, -0.001)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (follow-up 13	–91 days; unitless; l	petter indicated by	lower values)					
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	-0.027 (-0.033, -0.022)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol r	atio (follow-up 13–	91 days; unitless; be	etter indicated by l	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.029 (-0.034, -0.024)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	r values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹⁰	None	2010	-0.002 (-0.005, 0.002)	⊕⊕⊕O MODERATE	IMPORTANT
ApoA-I (follow-up	14–91 days; units	mg/dL per 1% ener	gy exchange; better	indicated by highe	ervalues)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹⁰	None	1425	-1.8 (-3.7, 0.1)	⊕⊕⊕O MODERATE	IMPORTANT
Apo-B (follow-up	14–91 days; units r	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-7.8 (-9.5, -6.0)	⊕⊕⊕⊕ HIGH	IMPORTANT

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; CI, confidence interval; LDL, low-density lipoprotein; HDL, high-density lipoprotein; RCT, randomized controlled trial

¹ A mixture of dietary saturated fatty acids (SFA) is replaced with *cis*-monounsaturated fatty acids (*cis*-MUFA) consisting primarily of oleic acid.

² Number of data points (diet groups) are provided in parentheses. Each data point contains mean dietary information on SFA, cis-polyunsaturated fatty acids (cis-PUFA) and cis-MUFA intake as well as an associated change in a given serum lipid or lipoprotein for each study group (i.e. intervention and control groups) at the end of a dietary treatment period, and was extracted for all treatment groups within studies included in the analysis.

³ All studies were strictly controlled, dietary trials lasting from 13 to 91 days in which protein and cholesterol intakes were held constant. Some of the studies with parallel design were assessed as having unclear risk of bias in terms of randomization as the randomization procedure was not described adequately. Studies with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have an significant, systematic effect on study results. Blinding was not deemed to be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some studies may have been able to distinguish between intervention and control diets, this was not

expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were judged as having a low risk of bias. The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of *cis*-MUFA.

- This analysis was conducted as a multiple regression in which data points (see Footnote 2) were directly extracted from each study, rather than extraction of mean differences between groups within each included study. As a result, directly measuring between-study variability in a quantitative manner is not feasible. Qualitative assessment of the included studies and the strength and consistency of the results of the multiple regression, however, indicate that any inconsistency is unlikely to decrease confidence in the results of the multiple regression analysis and is therefore not considered to be serious.
- 5 All studies directly assessed the effect of modifying dietary fat intake such that models could be derived, which provide estimates of the effects of exchanging SFA with other nutrients on serum lipids and lipoproteins, which were priority health outcomes decided upon prior to initiating review. All studies were conducted in the population of interest (adults without disturbances of lipid metabolism or diabetes).
- 6 Imprecision was assessed using the 95% CI of the regression coefficient as a proxy for the 95% CI of a pooled estimate of effect, the rationale being that the regression coefficient is a direct measure of the effect of reducing SFA intake on a particular serum lipid or lipoprotein and the 95% CI is a measure of variability of that effect. Unless otherwise noted, the 95% CI does not cross a threshold of irrelevant benefit or important harm and the outcome has not been downgraded for serious imprecision.
- ⁷ Publication bias was not formally assessed.
- ⁸ Total number of participants. Data points containing dietary information on SFA, *cis*-PUFA and *cis*-MUFA intake as well as an associated change in a given serum lipid or lipoprotein were directly extracted from all study groups from included trials without distinction between intervention and control groups.
- ⁹ The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of cis-MUFA.
- 10 The 95% CI crosses a threshold of important benefit or harm and the outcome has therefore been downgraded for serious imprecision.

GRADE evidence profile 3

Question: What is the effect of replacing saturated fatty acids in the diet of adults with carbohydrates?¹

Population: General adult population

			Quality assessment	:			No of	F#4-449		
No. of studies ²	Design	Risk of bias ³	Inconsistency ⁴	Indirectness ⁵	Imprecision ⁶	Other considerations ⁷	No. of participants ⁸	Effect ⁹ (95% CI)	Quality	Importance
Total cholesterol	(units mmol/L per	1% energy exchang	ge; better indicated	by lower values)						
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.041 (-0.047, -0.035)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (u	ınits mmol/L per 1º	% energy exchange	; better indicated b	y lower values)						
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.033 (-0.039, -0.027)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (units mmol/L per 1	% energy exchange	e; better indicated b	y higher values)						
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.010 (-0.012, -0.008)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (units	s mmol/L per 1% er	nergy exchange; be	tter indicated by lo	wer values)						
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	0.011 (0.007, 0.014)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (unitless; bet	ter indicated by low	ver values)						
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious Imprecision ¹⁰	None	1990	0.001 (-0.006, 0.007)	⊕⊕⊕O MODERATE	IMPORTANT
LDL cholesterol to	HDL cholesterol r	atio (unitless; bette	er indicated by lowe	er values)						
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.007 (-0.013, -0.001)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(unitless; better in	dicated by lower va	lues)						
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.014 (0.010, 0.018)	⊕⊕⊕⊕ HIGH	IMPORTANT
ApoA-I (units mg/	dL per 1% energy e	exchange; better in	dicated by higher va	alues)						
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	-7.0 (-9.0, -0.51)	⊕⊕⊕⊕ HIGH	IMPORTANT
Apo-B (units mg/d	dL per 1% energy e	xchange; better inc	licated by lower val	ues)						
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-3.6 (-5.4, -1.7)	⊕⊕⊕⊕ HIGH	IMPORTANT

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; CI, confidence interval; LDL, low-density lipoprotein; HDL, high-density lipoprotein; RCT, randomized controlled trial

A mixture of dietary saturated fatty acids (SFA) was replaced with a mixture of carbohydrates (mono-, di- and polysaccharides), however, the number of studies providing sufficient dietary information to determine, exactly, the composition of the carbohydrate used in the studies was limited.

² Number of data points (diet groups) are provided in parentheses. Each data point contains mean dietary information on SFA, *cis*-polyunsaturated fatty acid (*cis*-PUFA) and *cis*-monounsaturated fatty acid (*cis*-MUFA) intake as well as an associated change in a given serum lipid or lipoprotein for each study group (i.e. intervention and control groups) at the end of a dietary treatment period, and was extracted for all treatment groups within studies included in the analysis.

³ All studies were strictly controlled, dietary trials lasting from 13 to 91 days in which protein and cholesterol intakes were held constant. Some of the studies with parallel design were assessed as having unclear risk of bias in terms of randomization as the randomization procedure was not described adequately. Studies with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were randomized, as all participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have an significant, systematic effect on study results. Blinding was not deemed to

be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some studies may have been able to distinguish between intervention and control diets, this was not expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were judged as having a low risk of bias. The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of carbohydrates.

- This analysis was conducted as a multiple regression in which data points (see Footnote 2) were directly extracted from each study, rather than extraction of mean differences between groups within each included study. As a result, directly measuring between-study variability in a quantitative manner is not feasible. Qualitative assessment of the included studies and the strength and consistency of the results of the multiple regression, however, indicate that any inconsistency is unlikely to decrease confidence in the results of the multiple regression analysis and is therefore not considered to be serious.
- 5 All studies directly assessed the effect of modifying dietary fat intake such that models could be derived, which provide estimates of the effects of exchanging SFA with other nutrients on serum lipids and lipoproteins, which were priority health outcomes decided upon prior to initiating review. All studies were conducted in the population of interest (adults without disturbances of lipid metabolism or diabetes).
- ⁶ Imprecision was assessed using the 95% CI of the regression coefficient as a proxy for the 95% CI of a pooled estimate of effect, the rationale being that the regression coefficient is a direct measure of the effect of reducing SFA intake on a particular serum lipid or lipoprotein and the 95% CI is a measure of variability of that effect. Unless otherwise noted, the 95% CI does not cross a threshold of irrelevant benefit or important harm and the outcome has not been downgraded for serious imprecision.
- ⁷ Publication bias was not formally assessed.
- Total number of participants. Data points containing dietary information on SFA, cis-PUFA and cis-MUFA intake as well as an associated change in a given serum lipid or lipoprotein were directly extracted from all study groups from included trials without distinction between intervention and control groups.
- ⁹ The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of carbohydrates.
- 10 Imprecision was assessed as indicated in Footnote 6. The 95% CI crosses a threshold of important benefit or harm and the outcome has therefore been downgraded for serious imprecision.

GRADE evidence profile 4

Question: What is the effect of a reduction in saturated fatty acids intake in adults with intakes greater than 10% of total energy intake?¹

Population: General adult population

			Quality assessment				No. of	Effect ⁹		
No. of studies ²	Design	Risk of bias ³	Inconsistency⁴	Indirectness ⁵	Imprecision ⁶	Other considerations7	participants ⁸	(95% CI)	Quality	Importance
By replacing SFA	specifically with	cis-PUFA?10								
Total cholesterol	(follow-up 13–91 d	ays; units mmol/L p	oer 1% energy exch	ange; better indica	ted by lower values)				
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.064 (-0.070, -0.058)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	er 1% energy exchai	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.055 (-0.061, -0.050)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (follow-up 13–91 da	ays; units mmol/L po	er 1% energy excha	nge; better indicate	ed by higher values)				
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.005 (-0.006, -0.003)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; เ	units mmol/L per 1%	6 energy exchange;	better indicated by	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.010 (-0.014, -0.007)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (follow-up 13	–91 days; unitless; l	petter indicated by	lower values)			,		
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	-0.034 (-0.040, -0.028)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol r	atio (follow-up 13–	91 days; unitless; be	etter indicated by lo	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.034 (-0.040, -0.029)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	r values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.005 (-0.009, -0.002)	⊕⊕⊕⊕ HIGH	IMPORTANT
ApoA-I (follow-up	14–91 days; units	mg/dL per 1% energ	gy exchange; better	indicated by highe	er values)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	-4.9 (-7.3, -2.5)	⊕⊕⊕⊕ HIGH	IMPORTANT
Apo-B (follow-up	14–91 days; units r	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-10.2 (-12.4, -8.1)	⊕⊕⊕⊕ HIGH	IMPORTANT

By replacing SFA	A specifically with	cis-MUFA? ¹¹								
Total cholesterol	(follow-up 13–91 d	ays; units mmol/L բ	per 1% energy exch	ange; better indica	ted by lower values)				
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.046 (-0.051, -0.040)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	er 1% energy excha	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.042 (-0.047, -0.037)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (follow-up 13–91 da	ys; units mmol/L p	er 1% energy excha	nge; better indicat	ed by higher values))				
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.002 (-0.004, 0.000)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; ι	units mmol/L per 19	% energy exchange;	better indicated b	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.004 (-0.007, -0.001)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (follow-up 13	-91 days; unitless; l	better indicated by	lower values)					
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	-0.027 (-0.033, -0.022)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	o HDL cholesterol r	atio (follow-up 13–	91 days; unitless; b	etter indicated by l	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.029 (-0.034, -0.024)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD)L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	ervalues)			,		
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	2010	-0.002 (-0.005, 0.002)	⊕⊕⊕O MODERATE	IMPORTANT
ApoA-I (follow-up	14–91 days; units	mg/dL per 1% ener	gy exchange; better	indicated by highe	er values)			,		1
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	1425	-1.8 (-3.7, 0.1)	⊕⊕⊕O MODERATE	IMPORTANT
Apo-B (follow-up	14–91 days; units r	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-7.8 (-9.5, -6.0)	⊕⊕⊕⊕ HIGH	IMPORTANT
By replacing SF/	A specifically with	carbohydrates? ¹³	•		•					
Total cholesterol	(follow-up 13–91 d	ays; units mmol/L p	per 1% energy exch	ange; better indica	ted by lower values)				
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.041 (-0.047, -0.035)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (u	units mmol/L per 1	% energy exchange	; better indicated b	y lower values)						
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.033 (-0.039, -0.027)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (units mmol/L per 1	% energy exchange	e; better indicated b	y higher values)						
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.010 (-0.012, -0.008)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (unit	s mmol/L per 1% er	nergy exchange; be	tter indicated by lo	wer values)	•			·		
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	0.011 (0.007, 0.014)	⊕⊕⊕⊕ HIGH	IMPORTANT

			Quality assessment				No. of	Effect ⁹		Importance		
No. of studies ²	Design	Risk of bias ³	Inconsistency ⁴	Indirectness ⁵	Imprecision ⁶	Other considerations7	participants ⁸	(95% CI)	Quality			
Total cholesterol t	otal cholesterol to HDL cholesterol ratio (unitless; better indicated by lower values)											
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	1990	0.001 (-0.006, 0.007)	⊕⊕⊕O MODERATE	IMPORTANT		
LDL cholesterol to	HDL cholesterol r	atio (unitless; bette	rindicated by lowe	rvalues)								
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.007 (-0.013, -0.001)	⊕⊕⊕⊕ HIGH	IMPORTANT		
Triglyceride to HD	L cholesterol ratio	(unitless; better in	dicated by lower va	lues)								
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.014 (0.010, 0.018)	⊕⊕⊕⊕ HIGH	IMPORTANT		
ApoA-I (units mg/	dL per 1% energy e	xchange; better inc	licated by higher va	alues)								
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	-7.0 (-9.0, -0.51)	⊕⊕⊕⊕ HIGH	IMPORTANT		
Apo-B (units mg/c	IL per 1% energy ex	xchange; better ind	icated by lower val	ues)								
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-3.6 (-5.4, -1.7)	⊕⊕⊕⊕ HIGH	IMPORTANT		

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; CI, confidence interval; cis-MUFA, cis-monounsaturated fatty acids; cis-PUFA, cis-polyunsaturated fatty acids; LDL, low-density lipoprotein; HDL, high-density lipoprotein; RCT, randomized controlled trial; SFA, saturated fatty acids

- 1 Effects of decreasing SFA intake on serum lipids and lipoproteins by replacement with *cis*-PUFA, *cis*-MUFA or carbohydrates were observed across a wide range of saturated fatty acids intake, from 1.6 to 24.4% of total energy intake and residuals analysis indicates that the relationship between SFA intake and effect on serum lipids and lipoproteins is consistent across the entire range of SFA intakes. Of the 177 data points used in the multiple regression, 61 included an SFA intake component of more than 10% of total energy intake.
- ² Number of data points (diet groups) are provided in parentheses. Each data point contains mean dietary information on SFA, *cis*-MUFA and *cis*-PUFA intake as well as an associated change in a given serum lipid or lipoprotein for each study group (i.e. intervention and control groups) at the end of a dietary treatment period, and was extracted for all treatment groups within studies included in the analysis.
- 3 All studies were strictly controlled, dietary trials lasting from 13 to 91 days in which protein and cholesterol intakes were held constant. Some of the studies with parallel design were assessed as having unclear risk of bias in terms of randomization as the randomization procedure was not described adequately. Studies with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have an significant, systematic effect on study results. Blinding was not deemed to be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some studies may have been able to distinguish between intervention and control diets, this was not expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were judged as having a low risk of bias. The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of *cis*-PUFA, *cis*-MUFA or carbohydrates.
- This analysis was conducted as a multiple regression in which data points (see Footnote 2) were directly extracted from each study, rather than extraction of mean differences between groups within each included study. As a result, directly measuring between-study variability in a quantitative manner is not feasible. Qualitative assessment of the included studies and the strength and consistency of the results of the multiple regression, however, indicate that any inconsistency is unlikely to decrease confidence in the results of the multiple regression analysis and is therefore not considered to be serious.
- 5 All studies directly assessed the effect of modifying dietary fat intake such that models could be derived, which provide estimates of the effects of exchanging SFA with other nutrients on serum lipids and lipoproteins, which were priority health outcomes decided upon prior to initiating review. All studies were conducted in the population of interest (adults without disturbances of lipid metabolism or diabetes).
- ⁶ Imprecision was assessed using the 95% CI of the regression coefficient as a proxy for the 95% CI of a pooled estimate of effect, the rationale being that the regression coefficient is a direct measure of the effect of reducing SFA intake on a particular serum lipid or lipoprotein and the 95% CI is a measure of variability of that effect. Unless otherwise noted, the 95% CI does not cross a threshold of irrelevant benefit or important harm and the outcome has not been downgraded for serious imprecision.
- Publication bias was not formally assessed.
- Total number of participants. Data points containing dietary information on SFA, cis-PUFA and cis-MUFA intake as well as an associated change in a given serum lipid or lipoprotein were directly extracted from all study groups from included trials without distinction between intervention and control groups.
- The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of cis-PUFA, cis-MUFA or carbohydrates as indicated in the blue subheadings.
- 10 The *cis*-PUFA used as isocaloric replacement for SFA in individual studies assessing serum lipid and lipoprotein outcomes were linoleic acid and α -linolenic acid.
- 11 The cis-MUFA used as isocaloric replacement for SFA in individual studies assessing serum lipid and lipoprotein outcomes was predominantly oleic acid.
- 12 The 95% CI crosses a threshold of important benefit or harm and the outcome has therefore been downgraded for serious imprecision.
- ¹³ The carbohydrates used as isocaloric replacement for SFA in individual studies assessing serum lipid outcomes were a mixture of mono-, di- and polysaccharides, however, the number of studies providing sufficient dietary information to determine, exactly, the composition of the carbohydrates used in the studies was limited.

GRADE evidence profile 5

Question: What is the effect of a reduction in saturated fatty acids intake in adults to less than 10% of total energy intake?¹

Population: General adult population

			Quality assessment				No. of	Effect ⁹		
No. of studies ²	Design	Risk of bias ³	Inconsistency ⁴	Indirectness ⁵	Imprecision ⁶	Other considerations ⁷	participants ⁸	(95% CI)	Quality	Importance
By replacing SFA	specifically with	cis-PUFA?10								
Total cholesterol ((follow-up 13–91 da	ays; units mmol/L p	er 1% energy exch	ange; better indica	ted by lower values					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.064 (-0.070, -0.058)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (fo	ollow-up 13–91 day	ys; units mmol/L pe	er 1% energy exchar	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.055 (-0.061, -0.050)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L po	er 1% energy excha	nge; better indicate	ed by higher values)					
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.005 (-0.006, -0.003)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; u	nits mmol/L per 1%	6 energy exchange;	better indicated by	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.010 (-0.014, -0.007)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol t	to HDL cholesterol	ratio (follow-up 13	–91 days; unitless; b	petter indicated by	lower values)					
66 (159)5	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	-0.034 (-0.040, -0.028)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol ra	atio (follow-up 13–	91 days; unitless; be	etter indicated by lo	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.034 (-0.040, -0.029)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	r values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.005 (-0.009, -0.002)	⊕⊕⊕⊕ HIGH	IMPORTANT
ApoA-I (follow-up	14–91 days; units n	mg/dL per 1% energ	gy exchange; better	indicated by highe	er values)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	-4.9 (-7.3, -2.5)	⊕⊕⊕⊕ HIGH	IMPORTANT
Apo-B (follow-up	14–91 days; units m	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-10.2 (-12.4, -8.1)	⊕⊕⊕⊕ HIGH	IMPORTANT
By replacing SFA	specifically with	cis-MUFA? ¹¹								
Total cholesterol ((follow-up 13–91 da	ays; units mmol/L p	er 1% energy exch	ange; better indica	ted by lower values					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.046 (-0.051, -0.040)	⊕⊕⊕⊕ HIGH	IMPORTANT

			Quality assessment				No. of	Effect ⁹		
No. of studies ²	Design	Risk of bias ³	Inconsistency ⁴	Indirectness ⁵	Imprecision ⁶	Other considerations ⁷	participants ⁸	(95% CI)	Quality	Importance
LDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	r 1% energy excha	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.042 (-0.047, -0.037)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (follow-up 13–91 da	ys; units mmol/L pe	er 1% energy excha	nge; better indicat	ed by higher values)				
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.002 (-0.004, 0.000)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; ι	ınits mmol/L per 1%	6 energy exchange	better indicated b	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.004 (-0.007, -0.001)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (follow-up 13	-91 days; unitless;	petter indicated by	lower values)					
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	-0.027 (-0.033, -0.022)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol r	atio (follow-up 13–	91 days; unitless; b	etter indicated by l	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.029 (-0.034, -0.024)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	ervalues)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	2010	-0.002 (-0.005, 0.002)	⊕⊕⊕O MODERATE	IMPORTANT
ApoA-I (follow-up	14–91 days; units i	ng/dL per 1% energ	gy exchange; better	indicated by highe	ervalues)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	1425	-1.8 (-3.7, 0.1)	⊕⊕⊕O MODERATE	IMPORTANT
Apo-B (follow-up	14–91 days; units n	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-7.8 (-9.5, -6.0)	⊕⊕⊕⊕ HIGH	IMPORTANT
By replacing SFA	specifically with	carbohydrates? ¹³								
Total cholesterol	(follow-up 13–91 d	ays; units mmol/L p	er 1% energy exch	ange; better indica	ted by lower values)				
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	-0.041 (-0.047, -0.035)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (u	units mmol/L per 19	% energy exchange	; better indicated b	y lower values)						
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	-0.033 (-0.039, -0.027)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (units mmol/L per 1	% energy exchange	; better indicated b	y higher values)						
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	-0.010 (-0.012, -0.008)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (unit	s mmol/L per 1% er	nergy exchange; be	tter indicated by lo	wer values)						
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	0.011 (0.007, 0.014)	⊕⊕⊕⊕ HIGH	IMPORTANT

Total cholesterol	to HDL cholesterol	ratio (unitless; bett	ter indicated by low	vervalues)							
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	1990	0.001 (-0.006, 0.007)	⊕⊕⊕O MODERATE	IMPORTANT	
LDL cholesterol to	LDL cholesterol to HDL cholesterol ratio (unitless; better indicated by lower values)										
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.007 (-0.013, -0.001)	⊕⊕⊕⊕ HIGH	IMPORTANT	
Triglyceride to HD	Triglyceride to HDL cholesterol ratio (unitless; better indicated by lower values)										
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.014 (0.010, 0.018)	⊕⊕⊕⊕ HIGH	IMPORTANT	
ApoA-I (units mg/	dL per 1% energy e	xchange; better inc	dicated by higher va	alues)							
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	-7.0 (-9.0, -0.51)	⊕⊕⊕⊕ HIGH	IMPORTANT	
Apo-B (units mg/o	dL per 1% energy e	xchange; better ind	icated by lower val	ues)							
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	-3.6 (-5.4, -1.7)	⊕⊕⊕⊕ HIGH	IMPORTANT	

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; CI, confidence interval; cis-MUFA, cis-monounsaturated fatty acids; cis-PUFA, cis-polyunsaturated fatty acids; LDL, low-density lipoprotein; HDL, high-density lipoprotein; RCT, randomized controlled trial: SFA. saturated fatty acids

- ¹ Effects of decreasing SFA intake on serum lipids and lipoproteins by replacement with *cis*-PUFA, *cis*-MUFA or carbohydrates were observed across a wide range of saturated fatty acids intake, from 1.6 to 24.4% of total energy intake and residuals analysis indicates that the relationship between SFA intake and effect on serum lipids and lipoproteins is consistent across the entire range of SFA intakes. Of the 177 data points used in the multiple regression, 113 included an SFA intake component of less than 10% of total energy intake; 65 data points included intakes of less than 8%.
- ² Number of data points (diet groups) are provided in parentheses. Each data point contains mean dietary information on SFA, *cis*-MUFA and *cis*-PUFA intake as well as an associated change in a given serum lipid or lipoprotein for each study group (i.e. intervention and control groups) at the end of a dietary treatment period, and was extracted for all treatment groups within studies included in the analysis.
- 3 All studies were strictly controlled, dietary trials lasting from 13 to 91 days in which protein and cholesterol intakes were held constant. Some of the studies with parallel design were assessed as having unclear risk of bias in terms of randomization as the randomization procedure was not described adequately. Studies with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have an significant, systematic effect on study results. Blinding was not deemed to be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some studies may have been able to distinguish between intervention and control diets, this was not expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were judged as having a low risk of bias. The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of *cis*-PUFA, *cis*-MUFA or carbohydrates.
- ⁴ This analysis was conducted as a multiple regression in which data points (see Footnote 2) were directly extracted from each study, rather than extraction of mean differences between groups within each included study. As a result, directly measuring between-study variability in a quantitative manner is not feasible. Qualitative assessment of the included studies and the strength and consistency of the results of the multiple regression, however, indicate that any inconsistency is unlikely to decrease confidence in the results of the multiple regression analysis and is therefore not considered to be serious.
- 5 All studies directly assessed the effect of modifying dietary fat intake such that models could be derived, which provide estimates of the effects of exchanging SFA with other nutrients on serum lipids and lipoproteins, which were priority health outcomes decided upon prior to initiating review. All studies were conducted in the population of interest (adults without disturbances of lipid metabolism or diabetes).
- 6 Imprecision was assessed using the 95% CI of the regression coefficient as a proxy for the 95% CI of a pooled estimate of effect, the rationale being that the regression coefficient is a direct measure of the effect of reducing SFA intake on a particular serum lipid or lipoprotein and the 95% CI is a measure of variability of that effect. Unless otherwise noted, the 95% CI does not cross a threshold of irrelevant benefit or important harm and the outcome has not been downgraded for serious imprecision.
- Publication bias was not formally assessed.
- Total number of participants. Data points containing dietary information on SFA, cis-PUFA and cis-MUFA intake as well as an associated change in a given serum lipid or lipoprotein were directly extracted from all study groups from included trials without distinction between intervention and control groups.
- ⁹ The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of cis-PUFA, cis-MUFA or carbohydrates as indicated in the blue subheadings.
- 10 The cis-PUFA used as isocaloric replacement for SFA in individual studies assessing serum lipid and lipoprotein outcomes were linoleic acid and α -linolenic acid.
- 11 The cis-MUFA used as isocaloric replacement for SFA in individual studies assessing serum lipid and lipoprotein outcomes was predominantly oleic acid.
- ¹² The 95% CI crosses a threshold of important benefit or harm and the outcome has therefore been downgraded for serious imprecision.
- 13 The carbohydrates used as isocaloric replacement for SFA in individual studies assessing serum lipid outcomes were a mixture mono-, di- and polysaccharides, however, the number of studies providing sufficient dietary information to determine, exactly, the composition of the carbohydrates used in the studies was limited.

GRADE evidence profile 6

Question: What is the effect of an increase in saturated fatty acids intake in adults with a starting intake of less than 10% of total energy intake?¹

Population: General adult population

			Quality assessment				No. of	Effect ⁹		
No. of studies ²	Design	Risk of bias ³	Inconsistency⁴	Indirectness⁵	Imprecision ⁶	Other considerations ⁷	participants ⁸	(95% CI)	Quality	Importance
By replacing cis-	PUFA with SFA? ¹⁰									
Total cholesterol	(follow-up 13–91 da	ays; units mmol/L p	er 1% energy excha	ange; better indica	ted by lower values					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	0.066 (0.060, 0.073)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (f	ollow-up 13–91 day	ys; units mmol/L pe	r 1% energy exchar	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	0.058 (0.052, 0.064)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	er 1% energy excha	nge; better indicate	ed by higher values)					
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	0.005 (0.004, 0.007)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days; u	nits mmol/L per 1%	energy exchange;	better indicated by	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	0.010 (0.006, 0.014)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholesterol	ratio (follow-up 13	-91 days; unitless; b	petter indicated by	lower values)					
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	0.034 (0.027, 0.041)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol ra	atio (follow-up 13–	91 days; unitless; be	etter indicated by lo	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.035 (0.028, 0.041)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	r values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.004 (0.001, 0.008)	⊕⊕⊕⊕ HIGH	IMPORTANT
ApoA-I (follow-up	14–91 days; units r	mg/dL per 1% energ	y exchange; better	indicated by highe	rvalues)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	6.3 (3.9, 8.7)	⊕⊕⊕⊕ HIGH	IMPORTANT
Apo-B (follow-up	14–91 days; units m	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	10.3 (7.7, 12.8)	⊕⊕⊕⊕ HIGH	IMPORTANT
By replacing cis-	MUFA with SFA?11									
Total cholesterol	(follow-up 13–91 da	ays; units mmol/L p	er 1% energy excha	ange; better indica	ted by lower values					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	0.049 (0.043, 0.055)	⊕⊕⊕ HIGH	IMPORTANT

LDL cholesterol (f	ollow-up 13–91 da	ys; units mmol/L pe	er 1% energy exchai	nge; better indicate	ed by lower values)					
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	0.045 (0.039, 0.051)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (f	follow-up 13–91 da	ays; units mmol/L pe	er 1% energy excha	nge; better indicate	ed by higher values)					
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	0.003 (0.001, 0.004)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (follo	w-up 13–91 days;	units mmol/L per 1%	6 energy exchange;	better indicated b	y lower values)					
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	0.004 (0.000, 0.007)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholestero	l ratio (follow-up 13	–91 days; unitless; l	better indicated by	lower values)					
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1990	0.028 (0.021, 0.034)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol to	HDL cholesterol r	ratio (follow-up 13–	91 days; unitless; be	etter indicated by l	ower values)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.030 (0.024, 0.036)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride to HD	L cholesterol ratio	(follow-up 13–91 d	ays; unitless; bette	r indicated by lowe	rvalues)					
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	2010	0.001 (-0.003, 0.004)	⊕⊕⊕O MODERATE	IMPORTANT
ApoA-I (follow-up	14–91 days; units	mg/dL per 1% energ	gy exchange; better	indicated by highe	ervalues)					
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	2.7 (0.7, 4.8)	⊕⊕⊕⊕ HIGH	IMPORTANT
Apo-B (follow-up	14–91 days; units r	ng/dL per 1% energ	y exchange; better	indicated by lower	values)					
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	8.1 (6.1, 10.1)	⊕⊕⊕⊕ HIGH	IMPORTANT
By replacing car	bohydrates with	SFA? ¹³								•
Total cholesterol	(follow-up 13–91 d	lays; units mmol/L p	er 1% energy exch	ange; better indica	ted by lower values)					
74 (177)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2172	0.045 (0.038, 0.051)	⊕⊕⊕⊕ HIGH	IMPORTANT
LDL cholesterol (u	units mmol/L per 1	% energy exchange	; better indicated b	y lower values)						
69 (165)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2026	0.036 (0.030, 0.043)	⊕⊕⊕⊕ HIGH	CRITICAL
HDL cholesterol (units mmol/L per 1	.% energy exchange	; better indicated b	y higher values)						
68 (163)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2017	0.011 (0.010, 0.013)	⊕⊕⊕⊕ HIGH	IMPORTANT
Triglyceride (units	s mmol/L per 1% e	nergy exchange; be	tter indicated by lo	wer values)						
72 (172)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2156	-0.012 (-0.015, -0.008)	⊕⊕⊕⊕ HIGH	IMPORTANT
Total cholesterol	to HDL cholestero	l ratio (unitless; bett	ter indicated by low	ver values)						
66 (159)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	Serious imprecision ¹²	None	1990	-0.002 (-0.009, 0.005)	⊕⊕⊕O MODERATE	IMPORTANT

			Quality assessment				No. of	F#+9		Importance	
No. of studies ²	Design	Risk of bias ³	Inconsistency ⁴	Indirectness ⁵	Imprecision ⁶	Other considerations ⁷	participants ⁸	Effect ⁹ (95% CI)	Quality		
LDL cholesterol to	DL cholesterol to HDL cholesterol ratio (unitless; better indicated by lower values)										
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	0.007 (0.001, 0.014)	⊕⊕⊕⊕ HIGH	IMPORTANT	
Triglyceride to HD	Triglyceride to HDL cholesterol ratio (unitless; better indicated by lower values)										
67 (161)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	2010	-0.016 (-0.020, -0.012)	⊕⊕⊕⊕ HIGH	IMPORTANT	
ApoA-I (units mg/	dL per 1% energy e	xchange; better inc	dicated by higher va	alues)							
41 (102)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1425	8.4 (6.4, 10.5)	⊕⊕⊕⊕ HIGH	IMPORTANT	
Apo-B (units mg/c	IL per 1% energy ex	kchange; better ind	icated by lower val	ues)							
42 (104)	RCTs	No serious risk of bias	No serious inconsistency	No serious indirectness	No serious imprecision	None	1486	3.7 (1.7, 5.8)	⊕⊕⊕⊕ HIGH	IMPORTANT	

ApoA-I, apolipoprotein A-I; Apo-B, apolipoprotein B; CI, confidence interval; cis-MUFA, cis-monounsaturated fatty acids; cis-PUFA, cis-polyunsaturated fatty acids; LDL, low-density lipoprotein; HDL, high-density lipoprotein; RCT, randomized controlled trial: SFA. saturated fatty acids

- ¹ Effects of increasing SFA intake on serum lipids and lipoproteins by replacing *cis*-PUFA, *cis*-MUFA or carbohydrates with a mixture of SFA, were observed across a wide range of SFA intakes, from 1.6 to 24.4% of total energy intake and residuals analysis indicates that the relationship between SFA intake and effect on serum lipids and lipoproteins is consistent across the entire range of SFA intakes. Of the 177 data points used in the multiple regression, 113 included an SFA intake component of less than 10% of total energy intake; 65 data points included intakes of less than 8%.
- ² Number of data points (diet groups) are provided in parentheses. Each data point contains mean dietary information on SFA, *cis*-MUFA and *cis*-PUFA intake as well as an associated change in a given serum lipid or lipoprotein for each study group (i.e. intervention and control groups) at the end of a dietary treatment period, and was extracted for all treatment groups within studies included in the analysis.
- ³ All studies were strictly controlled, dietary trials lasting from 13 to 91 days in which protein and cholesterol intakes were held constant. Some of the studies with parallel design were assessed as having unclear risk of bias in terms of randomization as the randomization procedure was not described adequately. Studies with crossover and Latin square designs were deemed to be at low risk of bias for randomization whether or not it was specifically indicated if participants were intended to receive all treatments and thus it is unlikely that any differences at baseline would have an significant, systematic effect on study results. Blinding was not deemed to be a significant source of bias as all interventions consisted of food provision and though it is possible that participants in some studies may have been able to distinguish between intervention and control diets, this was not expected to alter compliance given the study design and conduct. All outcomes were objectively measured by chemical and mathematical means so risk of detection bias (i.e. bias resulting from non-blinded outcome assessment) was considered to be very low. There was no indication of widespread attrition bias or selective reporting and other sources of bias were minimal. Overall, the studies were judged as having a low risk of bias. The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as *cis*-PUFA, *cis*-MUFA or carbohydrates is replaced with an isocaloric amount of SFA.
- This analysis was conducted as a multiple regression in which data points (see Footnote 2) were directly extracted from each study, rather than extraction of mean differences between groups within each included study. As a result, directly measuring between-study variability in a quantitative manner is not feasible. Qualitative assessment of the included studies and the strength and consistency of the results of the multiple regression, however, indicate that any inconsistency is unlikely to decrease confidence in the results of the multiple regression analysis and is therefore not considered to be serious.
- 5 All studies directly assessed the effect of modifying dietary fat intake such that models could be derived, which provide estimates of the effects of exchanging SFA with other nutrients on serum lipids and lipoproteins, which were priority health outcomes decided upon prior to initiating review. All studies were conducted in the population of interest (adults without disturbances of lipid metabolism or diabetes).
- 6 Imprecision was assessed using the 95% CI of the regression coefficient as a proxy for the 95% CI of a pooled estimate of effect, the rationale being that the regression coefficient is a direct measure of the effect of reducing SFA intake on a particular serum lipid or lipoprotein and the 95% CI is a measure of variability of that effect. Unless otherwise noted, the 95% CI does not cross a threshold of irrelevant benefit or important harm and the outcome has not been downgraded for serious imprecision.
- ⁷ Publication bias was not formally assessed.
- Total number of participants. Data points containing dietary information on SFA, cis-PUFA and cis-MUFA intake as well as an associated change in a given serum lipid or lipoprotein were directly extracted from all study groups from included trials without distinction between intervention and control groups.
- The reported effect is the regression coefficient resulting from multiple regression. It is interpreted as the change in a particular serum lipid or lipoprotein when 1% of total energy intake as SFA is replaced with an isocaloric amount of cis-PUFA, cis-MUFA or carbohydrates as indicated in the blue subheadings.
- ¹⁰ The cis-PUFA being isocalorically exchanged with SFA in individual studies assessing serum lipid and lipoprotein outcomes were linoleic acid and -linolenic acid.
- 11 The cis-MUFA being isocalorically exchanged with SFA in individual studies assessing serum lipid and lipoprotein outcomes was predominantly oleic acid.
- 12 The 95% CI crosses a threshold of important benefit or harm and the outcome has therefore been downgraded for serious imprecision.
- 13 The carbohydrates being isocalorically exchanged with SFA in individual studies assessing serum lipid outcomes were a mixture mono-, di- and polysaccharides, however, the number of studies providing sufficient dietary information to determine, exactly, the composition of the carbohydrates used in the studies was limited.

References

- 1 WHO. WHO handbook for guideline development, 2nd edition. Geneva: World Health Organization (WHO); 2014 (http://apps.who.int/iris/bitstream/10665/145714/1/9789241548960_eng.pdf, accessed 15 August 2015).
- 2 Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P et al. GRADE: An emerging consensuson rating quality of evidence and strength of recommendations. BMJ. 2008;336 (7650):924–926 (http://www.ncbi.nlm.nih.gov/pubmed/18436948, accessed 01 October 2015).
- 3 Mensink RP, Zock PL, Kester AD, Katan MB. Effects of dietary fatty acids and carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials. Am J Clin Nutr. 2003;77(5):1146–1155 (http://www.ncbi.nlm.nih.gov/pubmed/12716665, accessed 15 November 2015).
- 4 Brussaard JH, Katan MB, Groot PH, Havekes LM, Hautvast JG. Serum lipoproteins of healthy persons fed a low-fat diet or a polyunsaturated fat diet for three months. A comparison of two cholesterol-lowering diets. Atherosclerosis. 1982;42(2–3):205–219 (http://www.ncbi.nlm.nih.gov/pubmed/7073802, accessed 01 October 2015).
- 5 Keys A, Anderson JT, Grande F. Prediction of serum-cholesterol responses of man to changes in fats in the diet. Lancet. 1957;273(7003):959–966 (http://www.ncbi.nlm.nih.gov/pubmed/13482259, accessed 01 October 2015).
- 6 Higgins JP, Green S. Handbook for systematic reviews of interventions. The Cochrane Collaboration. 2011 (http://handbook.cochrane.org/, accessed 01 October 2015).
- Taboratory Methods Committee of the Lipid Research Clinics P. Cholesterol and triglyceride concentrations in serum/plasma pairs. Clin Chem. 1977;23(1):60–63 (http://www.ncbi.nlm.nih.gov/pubmed/832373, accessed 15 November 2015).
- 8 Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. Clin Chem. 1972;18(6):499–502 (http://www.ncbi.nlm.nih.gov/pubmed/4337382, accessed 15 November 2015).
- 9 Grande F, Anderson JT, Keys A. Comparison of effects of palmitic and stearic acids in the diet on serum cholesterol in man. Am J Clin Nutr. 1970;23(9):1184–1193 (http://www.ncbi.nlm.nih.gov/pubmed/5450836, accessed 15 November 2015).
- 10 Hegsted DM, McGandy RB, Myers ML, Stare FJ. Quantitative effects of dietary fat on serum cholesterol in man. Am J Clin Nutr. 1965;17(5):281–295 (http://www.ncbi.nlm.nih.gov/pubmed/5846902, accessed 15 November 2015).
- 11 Diet,nutritionandthepreventionofchronicdiseases:reportofaJointWHO/FAOExpertConsultation. Geneva: World Health Organization, 2003. (http://www.who.int/nutrition/publications/obesity/WHO_TRS_916/en/, accessed 01 October 2015).
- 12 Goff LM, Cowland DE, Hooper L, Frost GS. Low glycaemic index diets and blood lipids: a systematic review and meta-analysis of randomised controlled trials. Nutr Metab Cardiovasc Dis. 2013;23(1):1–10 (http://www.ncbi.nlm.nih.gov/pubmed/22841185, accessed 2 November 2015).
- 13 Li Y, Hruby A, Bernstein AM, Ley SH, Wang DD, Chiuve SE, et al. Saturated Fats Compared With Unsaturated Fats and Sources of Carbohydrates in Relation to Risk of Coronary Heart Disease: A Prospective Cohort Study. J Am Coll Cardiol. 2015;66(14):1538-48. (http://www.ncbi.nlm.nih.gov/pubmed/26429077, accessed 2 November 2015).

- 14 Mensink RP, Katan MB. Effect of monounsaturated fatty acids versus complex carbohydrates on high-density lipoproteins in healthy men and women. Lancet. 1987;1(8525):122–125 (http://www.ncbi.nlm.nih.gov/pubmed/2879969, accessed 15 November 2015).
- 15 Mensink RP, de Groot MJM, van den Broeke LT, Severijnen-Nobels AP, Demacker PNM, Katan MB. Effects of monounsaturated fatty acids v complex carbohydrates on serum lipoproteins and apoproteins in healthy men and women. Metab. 1989;38(2):172–178 (http://www.ncbi.nlm.nih.gov/pubmed/2536459, accessed 15 November 2015).
- 16 Mattson FH, Grundy SM. Comparison of effects of dietary saturated, monounsaturated, and polyunsaturated fatty acids on plasma lipids and lipoproteins in man. J Lipid Res. 1985;26(2):194–202 (http://www.ncbi.nlm.nih.gov/pubmed/3989378, accessed 15 November 2015).
- 17 Grundy SM. Comparison of monounsaturated fatty acids and carbohydrates for lowering plasma cholesterol. N Engl J Med. 1986;314(12):745–748 (http://www.ncbi.nlm.nih.gov/pubmed/951504, accessed 15 November 2015).
- 18 Brussaard JH, Dallinga-Thie G, Groot PHE, Katan MB. Effects of amount and type of dietary fat on serum lipids, lipoproteins and apolipoproteins in man. A controlled 8-week trial. Arterioscler. 1980;36(4):515–527 (http://www.ncbi.nlm.nih.gov/pubmed/7417369, accessed 15 November 2015).
- 19 Brussaard JH, Katan MB, Groot PHE, Havekes LM, Hautvast JGAJ. Serum lipoproteins of healthy persons fed a low-fat diet or a polyunsaturated fat diet for three months. Atherosclerosis. 1982;42(2-3):205-219 (http://www.ncbi.nlm.nih.gov/pubmed/7073802, accessed 15 November 2015).
- 20 Mensink RP, Katan MB. Effect of a diet enriched with monounsaturated or polyunsaturated fatty acids on levels of low-density and high-density lipoprotein cholesterol in healthy women and men. N Engl J Med. 1989;321(7):436–441 (http://www.ncbi.nlm.nih.gov/pubmed/2761578, accessed 15 November 2015).
- 21 Harris WS, Connor WE, McMurry MP. The comparative reductions of the plasma lipids and lipoproteins by dietary polyunsaturated fats: salmon oil versus vegetable oils. Metab. 1983;32(2):179–184 (http://www.ncbi.nlm.nih.gov/pubmed/6827988, accessed 15 November 2015).
- 22 Becker N, Illingworth DR, Alaupovic P, Connor WE, Sundberg EE. Effects of saturated, monounsaturated, and omega-6 polyunsaturated fatty acids on plasma lipids, lipoproteins, and apoproteins in humans. Am J Clin Nutr. 1983;37(3):355–360 (http://www.ncbi.nlm.nih.gov/pubmed/6829481, accessed 15 November 2015).
- 23 Bonanome A, Grundy SM. Effect of dietary stearic acid on plasma cholesterol and lipoprotein levels. N Engl J Med. 1988;318(19):1244–1248 (http://www.ncbi.nlm.nih.gov/pubmed/3362176, accessed 15 November 2015).
- 24 Grundy SM, Nix D, Whelan MF, Franklin L. Comparison of three cholesterol lowering diets in normolipidemic men. JAMA. 1986;256(17):2351-2355 (http://www.ncbi.nlm.nih.gov/pubmed/773141, accessed 15 November 2015).
- 25 Katan MB, Berns MAM, Glatz JFC, Knuiman JT, Nobels A, de Vries JHM. Congruence of individual responsiveness to dietary cholesterol and to saturated fat in humans. J Lipid Res. 1988;29(7):883–892 (http://www.ncbi.nlm.nih.gov/pubmed/3411247, accessed 15 November 2015).
- 26 Grande F, Anderson JT, Keys A. Diets of different fatty acid composition producing identical serum cholesterol levels in man. Am J Clin Nutr. 1972;25(1):53–60 (http://www.ncbi.nlm.nih.gov/pubmed/5007597, accessed 15 November 2015).
- 27 Anderson JT, Grande F, Keys A. Independence of the effects of cholesterol and degree of saturation of the fat in the diet on serum cholesterol in man. Am J Clin Nutr. 1976;29(11):1184–1189 (http://www.ncbi.nlm.nih.gov/pubmed/998534, accessed 15 November 2015).
- 28 Wolf RN, Grundy SM. Influence of exchanging carbohydrate for saturated fatty acids on plasma lipids and lipoproteins in men. J Nutr. 1983;113(8):1521–1528 (http://www.ncbi.nlm.nih.gov/pubmed/6875693, accessed 15 November 2015).

- 29 Grundy SM, Florentin L, Nix D, Whelan MF. Comparison of monounsaturated fatty acids and carbohydratesforreducing raised levels of plasmacholesterol in man. Am J Clin Nutr. 1988;47(6):965–969 (http://www.ncbi.nlm.nih.gov/pubmed/3376911, accessed 15 November 2015).
- 30 Reiser R, Probstfield JL, Silvers A, Scott LW, Shorney ML, Wood RD et al. Plasma lipid and lipoprotein response of humans to beef fat, coconut oil and safflower oil. Am J Clin Nutr. 1985;42(2):190–197 (http://www.ncbi.nlm.nih.gov/pubmed/4025191, accessed 15 November 2015).
- 31 Laine DC, Snodgrass CM, Dawson EA, Ener MA, Kuba K, Frantz ID. Lightly hydrogenated soy oil versus other vegetable oils as a lipid-lowering dietary constituent. Am J Clin Nutr. 1982;35(4):683–690 (http://www.ncbi.nlm.nih.gov/pubmed/7200319, accessed 15 November 2015).
- 32 Lewis B, Hammett F, Katan M, Kay RM, Merkx I, Nobels A et al. Towards an improved lipid-lowering diet: additive effects of changes in nutrient intake. Lancet. 1981;2(8259):1310–1313 (http://www.ncbi.nlm.nih.gov/pubmed/6118716, accessed 15 November 2015).
- 33 Kay RM, Jacobs M, Katan MB, Lewis B. Relationship between changes in plasma lipoprotein concentrations and fecal steroid excretion in man during consumption of four experimental diets. Atherosclerosis. 1985;55(1):15-23 (http://www.ncbi.nlm.nih.gov/pubmed/2988576, accessed 15 November 2015).
- 34 McDonald BE, Gerrard JM, Bruce VM, Corner EJ. Comparison of the effect of canola oil and sunflower oil on plasma lipids and lipoproteins and on in vivo thromboxane A2 and prostacyclin production in healthy young men. Am J Clin Nutr. 1989;50(6):1382–1388 (http://www.ncbi.nlm.nih.gov/pubmed/2596428, accessed 15 November 2015).
- 35 Mensink RP, Katan MB. Effect of dietary transfatty acids on high-density and low-density lipoprotein cholesterol levels in healthy subjects. N Engl J Med. 1990;323(7):439–445 (http://www.ncbi.nlm.nih. gov/pubmed/2374566, accessed 01 October 2015) (http://www.ncbi.nlm.nih.gov/pubmed/2374566, accessed 15 November 2015).
- 36 Valsta LM, Jauhiainen M, Aro A, Katan MB, Mutanen M. Effects of a monounsaturated rapeseed oil and a polyunsaturated sunflower oil diet on lipoprotein levels in humans. Arterioscler Thromb. 1992;12(1):50–57 (http://www.ncbi.nlm.nih.gov/pubmed/1731858, accessed 15 November 2015).
- 37 Wahrburg U, Martin H, Sandkamp M, Schulte H, Assmann G. Comparative effects of a recommended lipid-lowering diet vs a diet rich in monounsaturated fatty acids on serum lipid profiles in healthy young adults. Am J Clin Nutr. 1992;56(4):678–683 (http://www.ncbi.nlm.nih.gov/pubmed/1414967, accessed 15 November 2015).
- 38 Zock PL, Katan MB. Hydrogenation alternatives: Effects of trans fatty acids and stearic acid versus linoleic acid on serum lipids and lipoproteins in humans. J Lipid Res. 1992;33(3):399–410 (http://www.ncbi.nlm.nih.gov/pubmed/1569387, accessed 01 October 2015).
- 39 Wardlaw GM, Snook JT. Effect of diets high in butter, corn oil, or high-oleic acid sunflower oil on serum lipids and apolipoproteins in men. Am J Clin Nutr. 1990;51(5):815–821 (http://www.ncbi.nlm. nih.gov/pubmed/2333839, accessed 15 November 2015).
- 40 Kwon JS, Snook JT, Wardlaw GM, and Hwang DH. Effects of diets high in saturated fatty acids, canola oil, or safflower oil on platelet function, thromboxane B2 formation, and fatty acid composition of platelet phospholipids. Am J Clin Nutr. 1991;54(2):351-358 (http://www.ncbi.nlm.nih.gov/pubmed/1677525, accessed 15 November 2015).
- 41 Ginsberg HN, Barr SL, Gilbert A, Karmally W, Deckelbaum R, Kaplan K et al. Reduction of plasma cholesterol levels in normal men on an American Heart Association Step 1 diet or a Step 1 diet with added monounsaturated fat. N Engl J Med. 1990;322(9):574–579 (http://www.ncbi.nlm.nih.gov/pubmed/2304504, accessed 15 November 2015).
- 42 Chan JK, Bruce VM, McDonald BE. Dietary alpha-linolenic acid is as effective as oleic acid and linoleic acid in lowering blood cholesterol in normolipidemic men. Am J Clin Nutr. 1991;53(5):1230–1234 (http://www.ncbi.nlm.nih.gov/pubmed/1673589, accessed 15 November 2015).

- 43 Wardlaw GM, Snook JT, Lin MC, Puangco MA, Kwon JS. Serum lipid and apolipoprotein concentrations in healthy men on diets enriched in either canola oil or safflower oil. Am J Clin Nutr. 1991;54(1):104–110 (http://www.ncbi.nlm.nih.gov/pubmed/1905474, accessed 15 November 2015).
- 44 Berry EM, Eisenberg S, Haratz D, Friedlander Y, Norman Y, Kaufmann NA et al. Effects of diets rich in monounsaturated fatty acids on plasma lipoproteins--the Jerusalem Nutrition Study: high MUFAs vs high PUFAs. Am J Clin Nutr. 1991;53(4):899–907 (http://www.ncbi.nlm.nih.gov/pubmed/2008870, accessed 15 November 2015).
- 45 Berry EM, Eisenberg S, Friedlander Y, Harats D, Kaufmann NA, Norman Y et al. Effects of diets rich in monounsaturated fatty acids on plasma lipoproteins--the Jerusalem Nutrition Study. II. Monounsaturated fatty acids vs carbohydrates. Am J Clin Nutr. 1992;56(2):394–403 (http://www.ncbi.nlm.nih.gov/pubmed/1636617, accessed 15 November 2015).
- 46 Kris-Etherton PM, Derr J, Mitchell DC, Mustad VA, Russell ME, McDonnell ET et al. The role of fatty acid saturation on plasma lipids, lipoproteins, and apolipoproteins: I. Effects of whole food diets high in cocoa butter, olive oil, soybean oil, dairy butter, and milk chocolate on the plasma lipids of young men. Metab. 1993;42(1):121–129 (http://www.ncbi.nlm.nih.gov/pubmed/8446039, accessed 15 November 2015).
- 47 Tholstrup T, Marckmann P, Jespersen J, Vessby B, Jart A, Sandstrom B. Effect on blood lipids, coagulation, and fibrinolysis of a fat high in myristic acid and a fat high in palmitic acid. Am J Clin Nutr. 1994;60(6):919–925 (http://www.ncbi.nlm.nih.gov/pubmed/7985634, accessed 15 November 2015).
- 48 Denke MA, Grundy SM. Comparison of effects of lauric acid and palmitic acid on plasma lipids and lipoproteins. Am J Clin Nutr. 1992;56(5):895–898 (http://www.ncbi.nlm.nih.gov/pubmed/1415008, accessed 15 November 2015).
- 49 Bonanome A, Pagnan A, Biffanti S, Opportuno A, Sorgato F, Dorella M et al. Effect of dietary monounsaturated and polyunsaturated fatty acids on the susceptibility of plasma low density lipoproteins to oxidative modification. Arterioscler Thromb. 1992;12(4):529–533 (http://www.ncbi.nlm.nih.gov/pubmed/1558840, accessed 15 November 2015).
- 50 Judd JT, Clevidence BA, Muesing RA, Wittes J, Sunkin ME, Podczasy JJ. Dietary trans fatty acids: Effects on plasma lipids and lipoproteins of healthy men and women. Am J Clin Nutr. 1994;59(4):861–868 (http://www.ncbi.nlm.nih.gov/pubmed/8147331, accessed 01 October 2015).
- 51 Sundram K, Hayes KC, Siru OH. Dietary palmitic acid results in lower serum cholesterol than does a lauric-myristic acid combination in normolipemic humans. Am J Clin Nutr. 1994;59(4):841–846 (http://www.ncbi.nlm.nih.gov/pubmed/8147328, accessed 15 November 2015).
- 52 Tholstrup T, Marckmann P, Jespersen J, Sandstrom B. Fat high in stearic acid favorably affects blood lipids and factor VII coagulant activity in comparison with fats high in palmitic and lauric acids. Am J Clin Nutr. 1994;59:371–377 (http://www.ncbi.nlm.nih.gov/pubmed/8310987, accessed 15 November 2015).
- 53 Zock PL, de Vries JHM, Katan MB. Impact of myristic acid versus palmitic acid on serum lipid and lipoprotein levels in healthy women and men. Arterioscler Thromb. 1994;14(4):567–575 (http://www.ncbi.nlm.nih.gov/pubmed/8148355, accessed 15 November 2015).
- 54 Barr SL, Ramakrishnan R, Johnson C, Holleran S, Dell RB, Ginsberg HN. Reducing total dietary fat without reducing saturated fatty acids does not significantly lower total plasma cholesterol concentrations in normal males. Am J Clin Nutr. 1992;55(3):675–681 (http://www.ncbi.nlm.nih.gov/pubmed/1550043, accessed 15 November 2015).
- 55 Ginsberg HN, Karmally W, Barr SL, Johnson C, Holleran S, Ramakrishnan R. Effects of increasing dietary polyunsaturated fatty acids within the guidelines of the AHA step 1 diet on plasma lipid and lipoprotein levels in normal males. Arterioscler Thromb. 1994;14(6):892–901 (http://www.ncbi.nlm. nih.gov/pubmed/8199179, accessed 15 November 2015).

- 56 Judd JT, Oh SY, Hennig B, Dupont J, Marshall MW. Effects of low fat diets differing in degree of fat unsaturation on plasma lipids, lipoproteins, and apolipoproteins in adult men. J Am Coll Nutr. 1988;7(3):223–234 (http://www.ncbi.nlm.nih.gov/pubmed/3392355, accessed 15 November 2015).
- 57 Marshall MW, Judd JT, Matusik EJ, Church J, Canary JJ. Effects of low fat diets varying in P/S ratio on nutrient intakes, fecal excretion, blood chemistry profiles, and fatty acids of adult men. J Am Coll Nutr. 1986;5(3):263–279 (http://www.ncbi.nlm.nih.gov/pubmed/3734273, accessed 15 November 2015).
- 58 Sundram K, Hayes KC, Siru OH. Both dietary 18:2 and 16:0 may be required to improve the serum LDL/HDL cholesterol ratio in normocholesterolemic men. J Nutr Biochem. 1995;6(4):179–187 (http://www.sciencedirect.com/science/article/pii/095528639500029Y, accessed 15 November 2015).
- 59 Iacono JM, Dougherty RM. Lack of effect of linoleic acid on the high-density-lipoprotein-cholesterol fraction of plasma lipoproteins. Am J Clin Nutr. 1991;53(3):660–664 (http://www.ncbi.nlm.nih.gov/pubmed/1900384, accessed 15 November 2015).
- 60 Lichtenstein AH, Ausman LM, Carrasco W, Jenner JL, Gualtieri LJ, Goldin BR et al. Effects of canola, corn, and olive oils on fasting and postprandial plasma lipoproteins in humans as part of a National Cholesterol Education Program Step 2 diet. Arterioscler Thromb. 1993;13(10):1533–1542 (http://www.ncbi.nlm.nih.gov/pubmed/8399091, accessed 15 November 2015).
- 61 LichtensteinAH, Ausman LM, Carrasco W, Jenner JL, Ordovas JM, Schaefer EJ. Hypercholesterolemic effect of dietary cholesterol in diets enriched in polyunsaturated and saturated fat. Dietary cholesterol, fat saturation, and plasma lipids. Arterioscler Thromb. 1994;14(1):168–175 (http://www.ncbi.nlm.nih.gov/pubmed/8274473, accessed 15 November 2015).
- 62 Lichtenstein AH, Ausman LM, Carrasco W, Gualtieri LJ, Jenner JL, Ordovas JM et al. Rice bran oil consumption and plasma lipid levels in moderately hypercholesterolemic humans. Arterioscler Thromb. 1994;14(4):549–556 (http://www.ncbi.nlm.nih.gov/pubmed/8148353, accessed 15 November 2015).
- 63 Dougherty RM, Allman MA, Iacono JM. Effects of diets containing high or low amounts of stearic acid on plasma lipoprotein fractions and fecal fatty acid excretion of men. Am J Clin Nutr. 1995;61(5):1120–1128 (http://www.ncbi.nlm.nih.gov/pubmed/7733038, accessed 15 November 2015).
- 64 Marckmann P, Sandstrom B, Jespersen J. Fasting blood coagulation and fibrinolysis of young adults unchanged by reduction in dietary fat content. Arterioscler Thromb. 1992;12(2):201–205 (http://www.ncbi.nlm.nih.gov/pubmed/1543694, accessed 15 November 2015).
- 65 Howard BV, Hannah JS, Heiser CC, Jablonski KA, Paidi MC, Alarif L et al. Polyunsaturated fatty acids result in greater cholesterol lowering and less triacylglycerol elevation than do monounsaturated fatty acids in a dose-response comparison in a multiracial study group. Am J Clin Nutr. 1995;62(2):392–402 (http://www.ncbi.nlm.nih.gov/pubmed/7625348, accessed 15 November 2015).
- 66 Fielding CJ, Havel RJ, Todd KM, Yeo KE, Schloetter MC, Weinberg V et al. Effects of dietary cholesterol and fat saturation on plasma lipoproteins in an ethnically diverse population of healthy young men. J Clin Invest. 1995;95(2):611–618 (http://www.ncbi.nlm.nih.gov/pubmed/7860745, accessed 15 November 2015).
- 67 Park S, Snook JT, Bricker L, Morroco M, Van Voorhis R, Stasny E et al. Relative effects of high saturated fatty acid levels in meat, dairy products, and tropical oils on serum lipoprotein and low-density lipoprotein degradation by mononuclear cells in healthy males. Metab. 1996;45(5):550–558 (http://www.ncbi.nlm.nih.gov/pubmed/8622596, accessed 15 November 2015).
- 68 Cater NB, Heller HJ, Denke MA. Comparison of the effects of medium-chain triacylglycerols, palm oil, and high oleic acid sunflower oil on plasma triacylglycerol fatty acids and lipid and lipoprotein concentrations in humans. Am J Clin Nutr. 1997;65(1):41–45 (http://www.ncbi.nlm.nih.gov/pubmed/8988911, accessed 15 November 2015).

- 69 Tholstrup T, Sandstrom B, Hermansen JE, Hølmer G. Effect of modified dairy fat on postprandial and fasting plasma lipids and lipoproteins in healthy young men. Lipids. 1998;33(1):11–21 (http://www.ncbi.nlm.nih.gov/pubmed/9470169, accessed 15 November 2015).
- 70 Mazier MJ, Jones PJH. Diet fat saturation and feeding state modulate rates of cholesterol synthesis in normolipidemic men. J Nutr. 1997;127(2):332–340 (http://www.ncbi.nlm.nih.gov/pubmed/9039836, accessed 15 November 2015).
- 71 Ginsberg HN, Kris-Etherton P, Dennis B, Elmer PJ, Ershow A, Lefevre M et al. Effects of reducing dietary saturated fatty acids on plasma lipids and lipoproteins in healthy subjects: the DELTA Study, protocol 1. Arterioscler Thromb Vasc Biol. 1998;18(3):441–449 (http://www.ncbi.nlm.nih.gov/pubmed/9514413, accessed 15 November 2015).
- 72 Müller H, Jordal O, Kierulf P, Kirkhus B, Pedersen JI. Replacement of partially hydrogenated soybean oil by palm oil in margarine without unfavorable effects on serum lipoproteins. Lipids. 1998;33(9):879–887 (http://www.ncbi.nlm.nih.gov/pubmed/9778135, accessed 01 October 2015).
- 73 Hunter KA, Crosbie LC, Weir A, Miller GJ, Dutta-Roy AK. A residential study comparing the effects of diets rich in stearic acid, oleic acid, and linoleic acid on fasting blood lipids, hemostatic variables and platelets in young healthy men. J Nutr Biochem. 2000;11(7-8):408–416 (http://www.ncbi.nlm. nih.gov/pubmed/11044636, accessed 15 November 2015).
- 74 Judd JT, Baer DJ, Clevidence BA, Kris-Etherton P, Muesing RA, Iwane M. Dietary cis and trans mono-unsaturated and saturated FA and plasma lipids and lipoproteins in men. Lipids. 2002;37(2):123–131 (http://www.ncbi.nlm.nih.gov/pubmed/11908904, accessed 01 October 2015).
- 75 Baer DJ, Judd JT, Clevidence BA, Tracy RP. Dietary fatty acids affect plasma markers of inflammation in healthy men fed controlled diets: a randomized crossover study. Am J Clin Nutr. 2004;79(6):969–973 (http://www.ncbi.nlm.nih.gov/pubmed/15159225, accessed 15 November 2015).
- 76 Vega-López S, Ausman LM, Jalbert SM, Erkkila AT, Lichtenstein AH. Palm and partially hydrogenated soybean oils adversely alter lipoprotein profiles compared with soybean and canola oils in moderately hyperlipidemic subjects. Am J Clin Nutr. 2006;84(1):54–62 (http://www.ncbi.nlm.nih. gov/pubmed/16825681, accessed 01 October 2015).
- 77 Lichtenstein AH, Ausman LM, Jalbert SM, Schaefer EJ. Effects of different forms of dietary hydrogenated fats on serum lipoprotein cholesterol levels. N Engl J Med. 1999;340(25):1933–1940 (http://www.ncbi.nlm.nih.gov/pubmed/10379016, accessed 01 October 2015).
- 78 Lovejoy JC, Smith SR, Champagne CM, Most MM, Lefevre M, DeLany JP et al. Effects of diets enriched in saturated (palmitic), monounsaturated (oleic), or trans (elaidic) fatty acids on insulin sensitivity and substrate oxidation in healthy adults. Diabetes Care. 2002;25(8):1283–1288 (http://www.ncbi.nlm.nih.gov/pubmed/12145222, accessed 01 October 2015).
- 79 Berglund L, Lefevre M, Ginsberg HN, Kris-Etherton PM, Elmer PJ, Stewart PW et al. Comparison of monounsaturated fat with carbohydrates as a replacement for saturated fat in subjects with a high metabolic risk profile: studies in the fasting and postprandial states. Am J Clin Nutr. 2007;86(6):1611–1620 (http://www.ncbi.nlm.nih.gov/pubmed/18065577, accessed 15 November 2015).
- 80 Binkoski AE, Kris-Etherton PM, Wilson TA, Mountain ML, Nicolosi RJ. Balance of unsaturated fatty acids is important to a cholesterol-lowering diet: comparison of mid-oleic sunflower oil and olive oil on cardiovascular disease risk factors. J Am Diet Assoc. 2005;105(7):1080–1086 (http://www.ncbi.nlm.nih.gov/pubmed/15983524, accessed 15 November 2015).
- 81 Castro P, Miranda JL, Gomez P, Escalante DM, Segura FL, Martin A et al. Comparison of an oleic acid enriched-diet vs NCEP-I diet on LDL susceptibility to oxidative modifications. Eur J Clin Nutr. 2000;54(1):61–67 (http://www.ncbi.nlm.nih.gov/pubmed/10694774, accessed 15 November 2015).
- 82 Kris-Etherton PM, Pearson TA, Wan Y, Hargrove RL, Moriarty K, Fishell V et al. High-monounsaturated fatty acid diets lower both plasma cholesterol and triacylglycerol concentrations. Am J Clin Nutr. 1999;70(6):1009–1015 (http://www.ncbi.nlm.nih.gov/pubmed/10584045, accessed 15 November 2015).

- 83 MüllerH, Lindman AS, Brantsaeter AL, Pedersen JI. The serum LDL/HDL cholesterol ratio is influenced more favorably by exchanging saturated with unsaturated fat than by reducing saturated fat in the diet of women. J Nutr. 2003;133(1):78–83 (http://www.ncbi.nlm.nih.gov/pubmed/12514271, accessed 15 November 2015).
- 84 Nielsen NS, Pedersen A, Sandstrom B, Marckmann P, Høy CE. Different effects of diets rich in olive oil, rapeseed oil and sunflower-seed oil on postprandial lipid and lipoprotein concentrations and on lipoprotein oxidation susceptibility. Br J Nutr. 2002;87(5):489–499 (http://www.ncbi.nlm.nih.gov/pubmed/12010587, accessed 15 November 2015).
- 85 Poppitt SD, Keogh GF, Mulvey TB, McArdle BH, MacGibbon AK, Cooper GJ. Lipid-lowering effects of a modified butter-fat: a controlled intervention trial in healthy men. Eur J Clin Nutr. 2002;56(1):64–71 (http://www.ncbi.nlm.nih.gov/pubmed/11840182, accessed 15 November 2015).
- 86 Rajaram S, Burke K, Connell B, Myint T, Sabate J. A monounsaturated fatty acid-rich pecan-enriched diet favorably alters the serum lipid profile of healthy men and women. J Nutr. 2001;131(9):2275–2279 (http://www.ncbi.nlm.nih.gov/pubmed/11533266, accessed 15 November 2015).
- 87 Sanders TA, Oakley FR, Crook D, Cooper JA, Miller GJ. High intakes of trans monounsaturated fatty acids taken for 2 weeks do not influence procoagulant and fibrinolytic risk markers for CHD in young healthy men. Br J Nutr. 2003;89(6):767–776 (http://www.ncbi.nlm.nih.gov/pubmed/12828793, accessed 01 October 2015).
- 88 Wagner KH, Tomasch R, Elmadfa I. Impact of diets containing corn oil or olive/sunflower oil mixture on the human plasma and lipoprotein lipid metabolism. Eur J Nutr. 2001;40(4):161–167 (http://www.ncbi.nlm.nih.gov/pubmed/11905957, accessed 15 November 2015).
- 89 Kratz M, Cullen P, Kannenberg F, Kassner A, Fobker M, Abuja PM et al. Effects of dietary fatty acids on the composition and oxidizability of low-density lipoprotein. Eur J Clin Nutr. 2002;56(1):72–81 (http://www.ncbi.nlm.nih.gov/pubmed/11840183, accessed 15 November 2015).
- 90 Lichtenstein AH, Matthan NR, Jalbert SM, Resteghini NA, Schaefer EJ, Ausman LM. Novel soybean oils with different fatty acid profiles alter cardiovascular disease risk factors in moderately hyperlipidemic subjects. Am J Clin Nutr. 2006;84(3):497–504 (http://www.ncbi.nlm.nih.gov/pubmed/16960162, accessed 01 October 2015).
- 91 Motard-Belanger A, Charest A, Grenier G, Paquin P, Chouinard Y, Lemieux S et al. Study of the effect of trans fatty acids from ruminants on blood lipids and other risk factors for cardiovascular disease. Am J Clin Nutr. 2008;87(3):593–599 (http://www.ncbi.nlm.nih.gov/pubmed/18326596, accessed 01 October 2015).
- 92 Rajaram S, Haddad EH, Mejia A, Sabate J. Walnuts and fatty fish influence different serum lipid fractions in normal to mildly hyperlipidemic individuals: a randomized controlled study. Am J Clin Nutr. 2009;89(5):1657S–1663S (http://www.ncbi.nlm.nih.gov/pubmed/19339404, accessed 15 November 2015).
- 93 Gillingham LG, Gustafson JA, Han SY, Jassal DS, Jones PJ. High-oleic rapeseed (canola) and flaxseed oils modulate serum lipids and inflammatory biomarkers in hypercholesterolaemic subjects. Br J Nutr. 2011;105(3):417–427 (http://www.ncbi.nlm.nih.gov/pubmed/20875216, accessed 01 October 2015).
- 94 Iggman D, Gustafsson IB, Berglund L, Vessby B, Marckmann P, Riserus U. Replacing dairy fat with rapeseed oil causes rapid improvement of hyperlipidaemia: a randomized controlled study. J Intern Med. 2011;270(4):356–364 (http://www.ncbi.nlm.nih.gov/pubmed/21466598, accessed 15 November 2015).
- 95 Marin C, Perez-Martinez P, Delgado-Lista J, Gomez P, Rodriguez F, Yubero-Serrano EM et al. The insulin sensitivity response is determined by the interaction between the G972R polymorphism of the insulin receptor substrate 1 gene and dietary fat. Mol Nutr Food Res. 2011;55(2):328–335 (http://www.ncbi.nlm.nih.gov/pubmed/20824664, accessed 15 November 2015).

- 96 Roussell MA, Hill AM, Gaugler TL, West SG, Heuvel JP, Alaupovic P et al. Beef in an Optimal Lean Diet study: effects on lipids, lipoproteins, and apolipoproteins. Am J Clin Nutr. 2012;95(1):9–16 (http://www.ncbi.nlm.nih.gov/pubmed/22170364, accessed 15 November 2015).
- 97 Zhao G, Etherton TD, Martin KR, West SG, Gillies PJ, Kris-Etherton PM. Dietary alpha-linolenic acid reduces inflammatory and lipid cardiovascular risk factors in hypercholesterolemic men and women. J Nutr. 2004;134(11):2991–2997 (http://www.ncbi.nlm.nih.gov/pubmed/15514264, accessed 15 November 2015).
- 98 Sabaté J, Haddad E, Tanzman JS, Jambazian P, Rajaram S. Serum lipid response to the graduated enrichment of a Step I diet with almonds: a randomized feeding trial. Am J Clin Nutr. 2003;77(6):1379–1384 (http://www.ncbi.nlm.nih.gov/pubmed/12791613, accessed 15 November 2015).
- 99 Curb JD, Wergowske G, Dobbs JC, Abbott RD, Huang B. Serum lipid effects of a high-monounsaturated fat diet based on macadamia nuts. Arch Intern Med. 2000;160(8):1154–1158 (http://www.ncbi.nlm. nih.gov/pubmed/10789609, accessed 15 November 2015).
- 100 Cater NB, Denke MA. Behenic acid is a cholesterol-raising saturated fatty acid in humans. Am J Clin Nutr. 2001;73(1):41–44 (http://www.ncbi.nlm.nih.gov/pubmed/11124748, accessed 15 November 2015).
- 101 Lacroix E, Charest A, Cyr A, Baril-Gravel L, Lebeuf Y, Paquin P et al. Randomized controlled study of the effect of a butter naturally enriched in trans fatty acids on blood lipids in healthy women. Am J Clin Nutr. 2012;95(2):318–325 (http://www.ncbi.nlm.nih.gov/pubmed/22205319, accessed 01 October 2015).

For further information please contact:

Nutrition Policy and Scientific Advice Unit (NPU)

Department of Nutrition for Health and Development (NHD)

World Health Organization (WHO)

20, Avenue Appia

CH–1211 Geneva 27

Switzerland

Fax: +41.22.791.4156 E-mail: NPUinfo@who.int

NHD website: http://www.who.int//nutrition







ISBN 978 92 4 156534 9

